



www.ajbrui.net

Afr. J. Biomed. Res. 14 (January 2011); 49 -55

Research article

Evaluation of Anti-diarrhoeal and Anti-ulcer Properties of Fractions of *Triumfetta cordifolia* A. Rich (Tiliaceae) Fruit in rats

***^a Nwafor, P. A., ^bEkpo, B.A.J., ^bAjibesin, K. A., ^bBala, D. N.**

^aDepartment of Pharmacology & Toxicology; ^bDepartment of Pharmacognosy and Natural Medicine, Faculty of Pharmacy, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.

ABSTRACT: *Triumfetta cordifolia* fruit is said to have many ethnobotanical uses, among them is their use in gastrointestinal disorders. The aim of the work is to investigate the antidiarrhoeal and antiulcerogenic properties of the fruit fractions in rats. The crude ethanol extract was dissolved in distilled water and partitioned with n-hexane, chloroform, ethyl acetate and n-butanol. 50 mg/kg of each of the fractions was investigated for antidiarrhoeal and antiulcerogenic effects in rats. Yohimbine, cimetidine and propranolol were used as positive controls. The fractions of the fruit, showed varying significant ($p < 0.001$), degrees of antidiarrhoeal and antiulcerogenic effects. It was observed that n-butanol and aqueous fractions had more effects on experimentally- induced ulcers and diarrhoea than chloroform and n-hexane fractions. The results on experimentally - induced ulcer and diarrhoeal models showed that the fractions possess these properties. These effects may in part be due to their actions on cholinergic and adrenergic mechanisms as well as their secondary metabolites.

Keywords: *Triumfetta cordifolia*, ulcer, diarrhoea, fractions, n-hexane, n-butanol, aqueous, chloroform.

INTRODUCTION

Triumfetta cordifolia A. Rich (Tiliaceae) is an erect, perennial herb with hairy but sometimes smooth stem commonly found in Florida, Bermuda, central and South America as well as in West Africa (Howard, 1989). In Nigeria, it is mainly found in the Southern region where it grows in sandy to clay soils with pH values between 5.5 to 8.0. In Akwa Ibom State, the root is used for venereal diseases, liver and kidney ailments while the fruit is usually macerated in water or local alcohol for the treatment of diarrhoea, dysentery,

ulcer and delayed labour (Idongesit Linus personal communication). The acute toxicity of extracts of this plant was earlier reported by Onilede (2005). To determine the medicinal properties of *Triumfetta cordifolia* A. Rich fruit as acclaimed by the local people of Uyo Local Government Area of Akwa Ibom State, the antidiarrhoeal and antiulcerogenic properties of the fruit were investigated on experimentally - induced diarrhoea and ulcer models in rats.

MATERIALS AND METHODS

Plant Material

The fruits of *Triumfetta cordifolia* were collected from Afaha Oku in Uyo Local Government Area of Akwa Ibom State in October, 2005. The plant was identified and authenticated by Dr. (Mrs.) Margaret Bassey of the Department of Botany, University of Uyo, where the specimen voucher was deposited.

Preparation of Extract

The fresh fruit (4 kg) was air-dried for four weeks after which it was reduced to powder using pestle and mortar. The pulverized plant material was then

*Address for correspondence:

paulnwafor@yahoo.com ; Tel: 234 803 677 8861;
234 805 915 3346

macerated in glass tank using 99% ethanol for 72h. The crude ethanolic extract was then concentrated to dryness in vacuo at 40°C. 100g of the ethanolic extract of the fruit was dissolved in 80 ml of distilled water and partitioned successive with n-hexane, chloroform, ethyl acetate and n-butanol. Partitioning with each of these solvents was done at least three times to ensure maximum separation (partitioning) of the constituents of the extract. The fractions were evaporated to dryness in vacuo at 40°C, monitored by TLC and subjected to pharmacological tests.

Phytochemical Screening

Phytochemical screening of the fruit was done according to the methods of Clarke (1975), Odebiyi and Sofowora (1978) and Trease and Evans (1989). Tests for alkaloids, saponins, tannins, flavonoids, cardiac glycosides, anthraquinones and terpenes were carried out.

Animals

Albino rats (weighing 140 – 200g) were used. They were housed in standardized environmental conditions (22 ± 2.5°C, relative humidity 80-85%, 12 h light / 12h dark cycle) and fed with standard rodent diet (Unfailing Veterinary Service, Uyo, Nigeria) and water ad libitum. Approval for the use of animals in the study was obtained from the Animal Ethics Committee, Faculty of Pharmacy, University of Uyo, Uyo, Akwa Ibom State, Nigeria.

Effect of *T. cordifolia* fruit fractions on intestinal propulsions in rats

The effects of *T. cordifolia* fruit fractions on intestinal propulsion in unanaesthetised rats was tested using the charcoal meal method of Capasso et al., (1976); Nwafor and Bassey (2007). They were fasted for 24h but allowed free access to water. They were randomized and placed in six cages of six animals per cage. Group 1 was administered with 10% Tween 80 (5 ml/kg) orally (p.o), using orogastric cannula. Groups 2-5 were pretreated with 50 mg/kg aqueous, n-butanol, chloroform and n-hexane fractions (p.o) respectively. Group 6 was pretreated with 5mg/kg diphenoxylate (p.o). After 1h, each rat was administered with 1ml charcoal meal (5 % activated charcoal suspended in 10 % aqueous tragacanth) orally. The rats were killed 30 min later by cervical dislocation and bled, and the small intestine rapidly dissected out and placed on a clean surface. The small intestine was carefully inspected and the distance traversed by the charcoal meal from the pylorus was measured. The length of the whole small intestine was also measured. The distance traversed by the charcoal meal from the pylorus was expressed as a

percentage of the distance from the pylorus to the ileocaecal junction.

Effect of *T. cordifolia* fruit fractions on castor oil – induced diarrhoea

Diarrhoea was induced in rats according to the method of Nwodo and Alumanah (1991); Nwafor and Bassey (2007). Animals were fasted for 24h but allowed free access to water. They were randomized into six groups of six rats each. Group 1 (control) received 10% Tween 80 (5 ml/kg) by gavage; groups 2-5 were treated with 50 mg/kg (p.o) of aqueous, n-butanol, chloroform and n-hexane fruit fractions of *Triumfetta cordifolia* respectively. Group 6 was administered with 5 mg/kg of yohimbine subcutaneously. After 1 h, each rat received 2 ml castor (p.o) and was then observed for consistency of faecal matter and the frequency of defaecation for 3 h.

Effect of *T. cordifolia* fruit fractions on castor oil – induced fluid accumulation

Fluid accumulation was induced in rats according to the method of DiCarlo et al., (1994); Nwafor and Bassey (2007). Animals were fasted for 24 h but allowed free access to water. They were randomized into six groups of six rats each. Group 1 (control) received 10 % Tween 80 (5 ml/kg) by gavage; groups 2-5 were given 50mg/kg of aqueous, n-butanol, chloroform and n-hexane fractions (p.o) respectively. Group 6 received yohimbine (5mg/kg.sc.). After 1 h, each rat received 2 ml of castor oil (p.o). Thirty minutes later, the rats were killed by cervical dislocation and exsanguinated, the small intestine was ligated at both pyloric sphincter and at the ileocaecal junctions. The entire small intestine was dissected out, its contents were expelled into a graduated measuring cylinder and the volume of the contents was recorded.

Effect of *T. cordifolia* fruit fractions on indomethacin –induced gastric ulceration in rats

The rats were randomized into six groups of six rats each. Food was withdrawn 24 h and water 2 h before the commencement of the experiment (Alphin and Ward, 1967). Group 1 (control) received only indomethacin (Sigma, 60 mg/kg p.o. dissolved in 5 % Na₂CO₃); groups 2-5 were pretreated with 50 mg/kg of aqueous, n-butanol, chloroform and n-hexane fractions of *Triumfetta cordifolia* (p.o.) respectively while Group 6 received cimetidine (100 mg/kg p.o.) 1 h later, group 2-6 were administered with indomethacin. 4 h after indomethacin administration, animals were killed by cervical dislocation. The stomachs were removed and opened along the greater curvature. The tissues were fixed with 10 % formaldehyde in saline. Macroscopic

examination was carried out with a hand lens and the presence of ulcer lesion was scored (Nwafor et al., 1996). Ulcer index (UI), preventive ratio (PR) and degree of ulceration (DU) of each of the groups pretreated with extract fraction was calculated using standard methods (Zaidi and Mukerji, 1958; Nwafor et al., 2000).

Effect of *T. cordifolia* fruit fractions on ethanol - induced gastric ulceration in rats

The above procedure (as in indomethacin - induced ulceration) was repeated using ethanol (2.5 ml/kg). The rats were randomized into six groups of six rats each. Food was withdrawn 24 h and water 2 h before the commencement of experiment. Group 1 (control) received only ethanol (2.5 ml/kg p.o.); groups 2-5 were pretreated with 50 mg/kg (p.o.) of aqueous, n-butanol, chloroform and n-hexane fruit fractions of *Triumfetta cordifolia* respectively; group 6 received propranolol (40 mg/kg; p.o) dissolved in distilled water). 1 h later, groups 2-5 were administered with ethanol. Four hours after ethanol administration, animals were killed by cervical dislocation. The stomachs were removed and opened along the greater curvature. The tissues were fixed with 10 % formaldehyde in saline. Macroscopic examination was carried out with a hand lens and the presence of lesions were scored (Nwafor et al., 2000).

Effect of *T. cordifolia* fruit fractions on reserpine - induced ulceration in rats

The above procedure (as in indomethacin - induced ulceration) was repeated using reserpine (8 mg/kg suspended in 12 % Tween 80). The rats were randomized into six groups of 6 rats each. Food was withdrawn 24 h and water 2 h before the commencement of experiment. Group 1 (control) received only reserpine (8 mg/kg p.o.); groups 2 – 5

were pretreated with 50 mg/kg of aqueous, n-butanol, chloroform and n-hexane fractions of *Triumfetta Cordifolia* (p.o) respectively. Group 6 received cimetidine (100 mg/kg p.o). 1 h later, groups 2 –5 were administered with reserpine. 18 h after administration of reserpine, animals were killed by cervical dislocation. The stomachs were removed and opened along the greater curvature. The tissues were fixed with 10 % formaldehyde in saline. Macroscopic examination was carried out with a hand lens and the presence of lesions were scored (Nwafor et al., 2005).

Statistical analysis

Multiple comparisons of Mean $\pm$ S.E.M were carried out by one way analysis of variance (ANOVA), followed by Tukey – Krammar multiple comparisons post tests at $p=0.05$.

RESULTS

Small intestinal propulsion

Different fractions of *Triumfetta cordifolia* inhibited the intestinal propulsion from 24.20 % to 48.2 %. The degree of inhibition was in this pattern: Aqueous > n-butanol > n-hexane > chloroform. These inhibitions were less than that exhibited by a standard drug (diphenoxylate 73.48 %) (Table 1).

Castor oil – induced diarrhoea

Extract fractions of *Triumfetta cordifolia* inhibited castor oil - induced diarrhoea in varying degrees ranging from n-butanol fraction (77.27 %) to chloroform fraction (40.91 %). The least inhibition exhibited by chloroform fraction was higher than that of yohimbine, an α_2 receptor blocker. (Table 2).

Table 1:

Effect of different fractions of *Triumfetta cordifolia* on intestinal propulsion in rats

Treatment/route of administration (p.o)	Dose (mg/kg)	Mean intestinal length (cm)	Mean distance moved by charcoal (cm)	% inhabitation
Control (10% Tween 80)	-	99.00 $\pm$ 3.19	63.30 $\pm$ 1.63	36.06
n-hexane	50	97.87 $\pm$ 5.54	60.02 $\pm$ 0.23 ^{ns}	38.67
Chloroform	50	103.47 $\pm$ 3.88	78.43 $\pm$ 0.27*	24.20
n-butanol	50	90.15 $\pm$ 9.49	48.52 $\pm$ 0.81*	46.18
Aqueous	50	91.32 $\pm$ 7.89	47.30 $\pm$ 0.75*	48.20
Diphenoxylate	5	94.27 $\pm$ 2.79	25.00 $\pm$ 0.57*	73.48

Values represent Mean $\pm$ S. E. M. (n=6) p.o = per oral

Significance relative to control: * $p<0.001$)

n.s = not significant

Table 2:Effect of different fractions of *Triumfetta cordifolia* on castor oil-induced diarrhoea in rats.

Treatment/route of administration (p.o)	Dose (mg/kg)	Mean faecal matter	% reduction
Control (10% Tween 80)	-	11.00+0.69	-
n-hexane	50	6.00+1.10*	45.45
Chloroform	50	6.50+0.47*	40.91
n-butanol	50	2.50 + 0.37*	77.27
Aqueous	50	5.67 + 0.61*	48.45
Yohimbine (sc)	5	7.83+0.67	31.82

Values represent Mean $\pm$ S. E. M. (n=6) p.o = per oral

Significance relative to control: *p<0.001)

n.s = not significant

Table 3:Effect of different fractions of *Triumfetta cordifolia* on castor oil-induced intestinal fluid accumulation in rats.

Treatment/route of administration (p.o)	Dose (mg/kg)	Mean volume of intestinal fluid (ml)	% reduction
Control (10% Tween 80)	-	3.32+0.10	-
n-hexane	50	2.92+0.01*	12.05
Chloroform	50	3.68+0.19*	-10.84
n-butanol	50	2.75 + 0.01*	17.17
Aqueous	50	2.80 + 0.01*	15.66
Yohimbine(Sc)	5	3.15+0.10ns	5.12

Values represent Mean $\pm$ S. E. M. (n =6), p.o= per oral; Sc = subcutaneous.

Significance relative to control: * P< 0.001;

Table 4:Effect of different fractions of *Triumfetta cordifolia* on indomethacin-induced ulceration in rats.

Treatment/route of administration (p.o)	Dose (mg/kg)	Degree of ulceration (%)	Ulcer index	Preventive ratio
Control (indomethacin)	60	100	10.75+0.03*	-
n-hexane	50	94.60	10.17+0.12*	5.40
Chloroform	50	78.30	8.42+0.06*	21.67
n-butanol	50	38.10	4.08+0.03*	62.05
Aqueous	50	60.40	6.39+0.14*	40.56
Cimetidine	100	1.42	0.95+0.03*	99.16

Values represent mean + S. E. M. (n=6); p.o., = per oral.

Significance relative to control: *P<0.001.

Castor oil – induced intestinal fluid accumulation

The effect of different fractions of *Triumfetta cordifolia* on castor oil-induced intestinal fluid accumulation is as shown on Table 3. The fractions showed minimal inhibitions on castor oil-induced fluid accumulation which ranged from chloroform (-10.84 %) to n-butanol (17.17 %). Relative to yohimbine, the fractions showed greater inhibitory effects.

Indomethacin - induced gastric ulceration

The fractions significantly (p<0.001) protected the animals from indomethacin- induced ulceration. This protection was in polarity-dependent fashion. Aqueous (93.50 %) > n-butanol (61.90 %) > chloroform(21.70 %) > n-hexane (5.40 %) (Table 4).

Ethanol induced gastric ulceration

Triumfetta cordifolia fractions pretreatment on ethanol-induced gastric ulceration, showed a non-polar dependent protection. This protection was statistically significant ($p < 0.001$) Table 5.

Table 5. Effect different fractions of *Triumfetta Cordifolia* on

Reserpine – induced gastric ulceration

The fractions significantly protected the rats from reserpine-induced ulceration in rats ($p < 0.001$). This protection is in non-polar dependent manner. (Table 6).

ethanol-induced ulceration in rats.

Treatment/route of administration (p.o)	Dose (mg/kg)	Degree of ulceration (%)	Ulcer index	Preventive ratio
Control (ethanol)	-	100	4.00+0.012*	-
n-hexane	50	75.00	3.00+0.02*	25.00
Chloroform	50	75.00	3.00+0.03*	25.00
n-butanol	50	37.50	1.50+0.02*	62.50
Aqueous	50	55.00	2.20+0.03*	45.00
Propranolol	40	62.50	2.50+0.50*	37.50

Values represent mean $\pm$ S. E. M. (n=6); p.o. =per oral.

Significance relative to control: * $P < 0.001$.

Table 6:

Effect of different fractions of *Triumfetta Cordifolia* in reserpine- induced ulceration in rats.

Treatment/route of administration (p.o)	Dose (mg/kg)	Degree of ulceration (%)	Ulcer index	Preventive ratio
Control (reserpine)	8	100.00	5.08+0.015	-
n-hexane	50	91.90	4.67+0.28 ^a	8.07
Chloroform	50	68.90	3.50+0.12 ^b	31.10
n-butanol	50	25.20	1.28+0.13 ^b	74.80
Aqueous	50	68.90	3.50+0.01 ^b	31.10
Cimetidine	100	90.20	4.58+0.22 ^b	9.84

Values represent Mean $\pm$ S. E. M. (n = 6); p.o = per oral.

Significance relative to control: ^a $p < 0.01$; ^b $p < 0.001$.

DISCUSSION

The extract fractions of *Triumfetta cordifolia* inhibited in non-uniform manner intestinal propulsion, castor oil - induced diarrhoea and fluid accumulation in rats. These degrees of inhibition were in this descending order: Aqueous > n-butanol > n-hexane > chloroform for intestinal propulsion; n-butanol > aqueous > n-hexane > chloroform for castor oil- induced diarrhea, and n – butanol > aqueous > n-hexane > chloroform for intestinal fluid accumulation respectively.

Spontaneous movement is inherent in non-vascular smooth muscle (myogenic) as a fundamental characteristics and is modified in - situ by hormonal and neuronal influences (Golenhofen, 1970; Nwafor et al., 2005). Spontaneous movement of gastrointestinal muscle is regulated by the cycles of depolarization and repolarization. It is known that acetylcholine binds to muscarinic receptors on smooth muscle, thereby triggering a sequence of events resulting ultimately in smooth muscle contraction (Nwafor and Okwuasaba, 2005). That the fractions had inhibitory effects on the spontaneous contractility of gastrointestinal muscle

indicated in part that it may be mediating its effect through cholinergic mechanism.

Drugs affecting motility, frequency and consistency of diarrhoea also affect secretion (DiCarlo et al., 1994; Nwafor and Okwuasaba, 2001). The intraluminal fluid accumulation induced by castor oil was enhanced by yohimbine, an α_2 - adrenergic blocker, suggesting the involvement of adrenergic mechanism.

All the results therefore suggest that the extract fractions produced an inhibitory action on gastrointestinal functions, motility and secretion and this is partly mediated by muscarinic and adrenergic receptor systems. The extract fractions also inhibited in varying degrees indomethacin, ethanol and reserpine-induced ulcerations in rats. The degrees of inhibition are in this descending order: n-butanol > aqueous > chloroform > n – hexane for indomethacin; n-butanol > aqueous > chloroform Ξ n-hexane for ethanol, and n-butanol > aqueous Ξ chloroform > n-butanol for reserpine respectively.

Indomethacin is a known ulcerogen especially in an empty stomach (Bhargava et al., 1973), the evidence of indomethacin - induced ulceration is mostly on the glandular (mucosal) part of the stomach (Evbunwa

and Bolarinwa, 1990; Nwafor and Bassey 2007). Although the mechanisms underlying the ulcerogenicity of indomethacin are not completely understood, it has been known that inhibition of prostaglandin synthesis may be important (Hiruma – Lima et al., 2006; Vane, 1971). This view is supported by the fact that prostaglandins normally serve as protective functions in stomach by maintaining gastric microcirculation (Vane, 1971; Ferreira and Vane, 1974) and causes gastric secretion of bicarbonate (Garner et al., 1979) and mucus (Menguy and Desbaillets, 1967). Numerous studies have demonstrated that the effect of absolute ethanol on the gastric mucosa is related to production of free radicals, the increase of lipoperoxidation and the decrease of the levels of non protein SH compounds in the gastric mucosa (Mizui and Doteuchi, 1986). It is also established that disturbances in gastric secretion, damage to the gastric mucosa, alterations in permeability, gastric mucus depletion and free radical production are observed after the administration of ethanol (Salim, 1990).

Reserpine is said to elucidate ulcers through the mobilization of superoxide and hydroxyl radicals (oxygen – derived free radicals (Salim, 1989). The fractions reduced reserpine-induced ulceration in rats. It has been proposed that mucosal protection induced by non - prostanoid compounds may be mediated through the mobilization of endogenous prostaglandins (Konturek et al., (1989). It is possible that one of the mechanisms of antiulcerogenic effects of the extract may be due to its ability to mobilize prostaglandins in gastric mucosa by increasing its microcirculation or through the inhibition of oxygen – derived free radicals formation in the gastric mucosa. All the results therefore suggest that the extract fractions produced inhibitory action on gastrointestinal functions, motility and secretion, and this effect is mediated in part through cholinergic and adrenergic mechanisms.

The phytochemical analysis of the extract fractions revealed the presence tannins, alkaloids, flavonoid and cardiac glycosides. Flavonoids and tannins are some of most important botanical compounds with anti-ulcer and gastroprotective activities (Wahida et al., 2007). Different mechanisms have been proposed to explain the gastroprotective effect of flavonoids, including an increase in mucosal prostaglandin content and a decrease in histamine secretion from mast cells by the inhibition of histidine decarboxylase. Flavonoids are free radical scavengers that are known to play an important role in preventing ulcerative and erosive lesions of gastrointestinal tract (Borrelli and Izzo, 2000; Konan and Bacchi, 2007) Tannins with its protein precipitating and vasoconstrictory effects could be advantageous in preventing ulcer development (Aguwa

and Nwako, 1988). Tannins being astringent, may have precipitated microproteins on site of the ulcer thereby forming an impervious protective pellicle over the lining to prevent absorption of toxic substances and resist the attack of proteolytic enzymes (Nwafor et al., 1996).

In conclusion therefore, the fractions may in part be mediating their actions through the cholinergic and adrenergic mechanisms or through the action of its active metabolites.

Acknowledgements

The authors gratefully acknowledge the technical assistance of Miss Sifonobong Akpan, Mr. Nsikan Udo and Mr. Aniefiok Ukpog, all of the Department of Pharmacology & Toxicology, University of Uyo, Uyo, Akwa Ibom State, Nigeria.

REFERENCES

- Aguwa, C. N., Nwako, S.O. (1988)** Preliminary Studies on the root extracts of *Nuclea latifolia* Smith, for Anti-ulcer Properties. *Nig. J. Pharm. Sc.* 4(1), 16 - 23.
- Alphin, R. S., Ward, J.W. (1967)** Action of Hexapyrronium bromide on gastric secretion in dogs and on gastric secretion and ulceration in rats. *Ach. Int. Pharm. Ther.* 270, 128 – 140.
- Bhargava, K. P., Gupta, M. B., Tangri, K. K. (1973)** Mechanism of ulcerogenic activity of indomethacin and oxyphenbutazone. *Eur. J. Pharmacol.* 22, 191-195.
- Borrelli, F. Izzo, A. A. (2000)** The plant Kingdom as a source of anti-ulcer remedies. *Phytother. Res.* 14, 581-591.
- Clarke, E. G. C. (Ed), (1975)** *Isolation and Identification of Drugs.* Vol. 2 Pharmaceutical Press, London. P. 905.
- Capasso, F., De Ruggiero, G., DiRosa, M., Sorrention L., (1976)** *Ricereche Farmacologiche Sulla 4-amino-5-Cloro-4 (2-dietilaminetil) 2-metossi benzamide.* *Boll. Ch. Farm.* 115, 649-657.
- DiCarlo, G. D., Mascolo, N., Izzo, A.A., Capasso, F., Autore, G. (1994)** Effects of Quercetin on gastrointestinal tract in rats and mice. *Phytother. Res.* 8, 42-45
- Evbuonwa, M. J., Bolarinwa, A. F. (1990)** Effect of diet on indomethacin-induced peptic ulceration in pregnant rats. *Nig. J. Physiol. Sc.* 6, 189-191.
- Ferreira, S. J., Vane, J. R. (1974)** New aspects of the mode of actions of NSAIDS. *Ann. Rev. Pharmacol.* 14, 57-70.
- Garner, A., Flemstrom, G., Heylings, J. R. (1979).** Effects of anti-inflammatory agents and prostalands on acid and bicarbonate secretions in the amphibian isolated gastric mucosa. *Gastroent.* 77, 451-457.
- Golenhofen, K. (1970)** slows rhythms in smooth muscle (minute rhythms). In: Bulbring E., Brady A. F., Jones, A. W., Tomita T (eds.) *Smooth Muscle.* London Edward Arnold Publishers, P. 316-342.
- Hiruma-Lima, C. A., Calvo, T. R., Rodrigues, C. M., Andrade, F. D. P., Vilegas, W., Brito, A.M.S., (2006)** Antiulcerogenic activity of *Alchornea castaneafolia*:

- effects on somatostatin, gastrin and prostaglandin. J Ethnopharmacol 83, 55-60.
- Howard, R. A. (1989)** Flora of the lesser Antilles Leeward and Windward Islands (Vol. 5). Arnold Arboretum, Harvard University, Jamaica Plain M. A. p.604.
- Konan, N. A., Bacchi, E. M., (2007)** Antiulcerogenic effect and acute toxicity of a Hydroethanolic extract from the cashew (*Anacardium Occidentale* L.) Leaves. J. Ethnopharmacol. 112, 237-242.
- Konturek, S.J., Redeki, T., Piastucki, I., Brzozowski, T., Drozdowicz, D., (1987)** Gastroprotection by colloidal bismuth subcitrate (De-nol) and Sucralfate: Role of endogenous prostaglandins. *Gut* 228, 201-205.
- Menguy, R., Desbaillets, L., (1967)** Role of inhibition of gastric mucous secretion in the phenomenon of gastric mucosal injury by indomethacin. *Am. J. Dig. Dis.* 12, 862-866.
- Mizui, T., Dteuchi, M., (1986)** Lipid peroxidation: a possible role in gastric damage induced by ethanol in rats. *Life Sc.* 38, 2163-2167.
- Nwafor, P.A., Effraim, K. D., Jacks, T.W. (1996).** Gastroprotective effects of Aqueous extract of *Khaya senegalensis* bark on indomethacin-induced ulceration in rats W. Afr. J. Pharmacol. Drug Res. 12, 46-50.
- Nwafor, P.A., Okwuasaba, F.K., Binda, L.G. (2000)** Antidiarrhoeal and antiulcerogenic effects of methanolic extract of *Asparagus pubescens* roots in rats, J. Ethnopharmacol. 72, 421-427.
- Nwafor, P.A. Okwuasaba, F.K. (2001)** The effect of methanolic extract of *Cassia nigricans* leaves on gastrointestinal tract. *Fitoter.* 72, 206-214.
- Nwafor, P. A., Jacks, T. W., Ekanem, A. U., Ching, F. Poh., (2005)** Antiulcerogenic and antidiarrhoeal potentials of methanolic extract of *Pausinystalia macroceras* stem-bark in rats. *Nig. J. Nat. Prod. Med.* 9, 63 – 63-67.
- Paul A. Nwafor, Okwuasaba, F. K., Uguru, M. O. (2005)** Effect of methanolic extract of *Asparagus pubescens* root on non-vascular smooth muscle in experimental animals. W. Afr. J. Pharmacol. Drug Res. 21, 31 – 36.
- Nwafor, P. A., Augustine I. L. Bassey (2007)** Evaluation of anti-diarrhoeal and anti-ulcerogenic potential of ethanol extract of *Carpolobia lutea* leaves in rodents J. Ethnopharmacol. 111, 619 – 624.
- Nwodo, O. F. C., Alumanah, E. O. (1991).** Studies of *Abrus precatorius* seed II. Antidiarrhoeal activity. J Ethnopharmacol 31, 395 – 398.
- Odebiyi, O. O., Sofowora, E. A. (1978)** Phytochemical screening of Nigerian Medicinal plants. *Lloydia* 41, 234 – 235.
- Onilede, F. (2005)** Investigation on antimicrobial effects of *Triumfetta cordifolia* A. Rich (Tiliaceae). Dissertation submitted to Department of Pharmacognosy & Natural medicine for the award of Bachelor of Pharmacy degree, University of Uyo, Nigeria pp. 20-45.
- Salim, A. S., (1989).** Role of oxygen – derived free radicals in the mechanism of chronic gastric ulceration in rat: implications for cytoprotection. *Dig.* 1 – 2, 113 – 119.
- Salim, A. S. (1990)** Removing oxygen-derived free radicals stimulates healing of ethanol-induced erosive gastritis in the rats. *Dig.* 47, 24-28.
- Trease, G. E., Evans, W. C., (1989)** *Pharmacognosy*. English Language Book Society, 13th ed. Bailliere Tindall, London Pp683-684.
- Vane, J.R. (1971)** Inhibition of prostaglandin synthesis as a mechanism of action of aspirin like drugs. *Nat. New Biol.* 231, 232-235.
- Wahida, B., Abderrahman, B., Nabil, C. (2007)** Antiulcerogenic activity of *Zizyphus lotus* (L.) extracts. J. Ethnopharmacol. 112, 228-231.
- Zaidi, S. H., Mukerji, B. (1958)** Experimental Peptic ulceration. Part I. The significance of mucous barrier. *Ind. J. Med. Res.* 46, 27-37.