

**BIOCONTROL OF POTATO WILT BY SELECTIVE RHIZOSPHERIC AND
ENDOPHYTIC BACTERIA ASSOCIATED WITH POTATO PLANT**

**Rado R^{1*}, Andrianarisoa B², Ravelomanantsoa S³,
Rakotoarimanga N¹, Rahetlah V⁴, Fienena FR¹ and O Andriambeloson¹**



Rado Rasolomampiannina

*Corresponding author mail: mampionina@yahoo.fr

¹Centre National de Recherche sur l'Environnement (CNRE), Antananarivo, Madagascar

²Laboratoire de Biotechnologie-Microbiologie, Département de Biochimie Fondamentale et Appliquée, Faculté des Sciences, Université d'Antananarivo, Madagascar

³CIRAD-3P, Pôle de Protection des Plantes, UMR PVBMT, Saint-Pierre, La Réunion, France

⁴(FIFAMANOR) Centre de Développement Rural et de Recherche Appliquée, Antsirabe, Madagascar

ABSTRACT

Ralstonia solanacearum is the causative agent of wilt disease in plants, which constitutes a severe problem to agricultural crops, particularly for potato production in Madagascar. The present study focuses on the isolation, *in vitro* and *in vivo* assays of potential rhizospheric and endophytic bacteria associated with healthy potato plant, capable to inhibit the growth of *Ralstonia solanacearum* for controlling potato bacterial wilt. A total of 77 bacteria strains were isolated from six soil rhizospheric samples and six vegetal material samples of healthy potatoes in the district of Antsirabe II. Forty of them were telluric actinomycetes, 25 were endophytic actinomycetes and 12 were fluorescent *Pseudomonas* spp. An additional 30 phytopathogenic isolates were obtained from six rhizospheric soil samples of diseased potatoes. Morphological, cultural, biochemical characterization and molecular identification with the *Ralstonia solanacearum* specific primers 759/760 revealed that 24 of the pathogenic isolates belong to the *Ralstonia solanacearum* species, biovar two; the causal agent of potato bacterial wilt. Isolates from healthy plants were, then, examined *in vitro* and *in vivo* for their antagonistic activity against *Ralstonia solanacearum* strain for their potential to improve potato plant growth. *In vitro* antagonism of actinomycete and *Pseudomonas* isolates against *Ralstonia solanacearum* development was performed using agar diffusion technique, while *in vivo* tests were conducted under greenhouse conditions. Ten antagonistic strains including two *Pseudomonas*, four telluric actinomycetes, and four endophytic actinomycetes inhibited the tested *Ralstonia* strain. Four strains, E7, E13 (endophytic actinomycete from root potatoes), S25 (telluric actinomycetes) and P7 (fluorescent *Pseudomonas*), showed high antagonistic activity against the pathogen with zones of inhibition from 23 to 40 mm. Of the four strains tested in greenhouse, E7 significantly reduced ($p < 0.05$) the percentage of *Ralstonia solanacearum* that infected plants by 72.04%. The isolates E13 and S25 have also been demonstrated to improve plant growth by increase of plant height to 44.63% and 44.84%, fresh weight to 68.75% and 75.85% and dry weight to 86.17% and 115.42%, respectively compared with non-treated control. Morphological and cultural characterization of these three active isolates showed that they belong to the genus *Streptomyces*. The antagonism of these isolates against *Ralstonia solanacearum* according to *in vitro* and *in vivo* tests results, along with their high efficiency as regards the improvement of plant development, suggests that these three actinomycete strains E7, E13 and S25 could be useful for biocontrol of potato bacterial wilt.

Key words: Potato, *Ralstonia*, Actinomycetes, *Pseudomonas*, Biocontrol

INTRODUCTION

Bacterial wilt caused by the various subgroups of the plant pathogenic bacterium *Ralstonia solanacearum* is an important disease of crop plants in tropical, subtropical and warm temperate regions. This pathogen has a wide host range and infects more than 450 monocotyledonous and dicotyledonous plant species belonging to 54 botanical families [1].

Bacterial wilt of potato (*Solanum tuberosum* L.) is a serious constraint to potato production and exportation in small-scale farms in the Vakinankaratra region, the main potato-farming region in Madagascar. The disease occurs in almost all potato-growing areas including paddy rice fields and causes severe losses of potato yield and quality specifically during the warm and rainy season. Visible symptoms are foliage wilting and stunting which could be followed by tubers rotting showing a vascular ring or a grey-brown discoloration of vascular tissues.

Infested soil and infected potato seed tubers and plants are the main sources of disease inoculum but contaminated surface water used for irrigation and contaminated tools may also spread the disease. As *Ralstonia solanacearum* strains can survive in soil and in the rooting system of many hosts including weeds, soil fumigation and crop rotation alone cannot control bacterial wilt. The use of resistant varieties of potato has significantly reduced bacterial wilt incidence but resistance is strain-specific and varies with temperature, soil moisture and other by environmental conditions. Many studies report that biological control is a promising alternative strategy for the integrated control of bacterial plant diseases. Therefore, the effectiveness of microbial antagonists against *Ralstonia solanacearum* growth has been reported in the present work. Among these antagonistic microorganisms, actinomycetes are well-known and classified among the most active [2].

In addition, some *Pseudomonas* spp. have been shown to initiate the expression of plant defense mechanisms [3].

The bioprotection conferred by these microbial agents may be based on competition for essential nutrients, antagonistic activity against the growth of pathogens through the production of antibiotics or enzymes, or their ability to stimulate defense systems in the plant host. In the wild, the rhizosphere of healthy plants includes *Pseudomonas* and/or actinomycetes capable of ensuring the defense of the plants against bacterial agents and/or phytopathogenic fungi.

Thus, the objective of this study was to identify possible biocontrol strains against bacterial wilt by isolating bacteria (actinomycetes and *Pseudomonas* strains) from healthy potato plants, evaluating *in vitro* their potential antagonism against isolated and characterized *Ralstonia solanacearum* from infected soil, and testing their plant growth potential under greenhouse conditions.

MATERIALS AND METHODS

Soil and plant sampling

Soil samples were collected in the rhizosphere of two potato varieties (Meva and Diamondra) in two sites (Antanetibe and Ankotrabe ambony) of the district of Antsirabe II in Vakinankaratra region, the main production region of potato. For telluric strains isolation, soil samples were taken under healthy potato plants. For endophytic actinomycetes isolation, three samples of healthy plant material (leaves, stems and roots) were used, and for phytopathogenic bacteria isolation, soil samples were collected under diseased potato plants.

Twelve soil samples were collected from these two sites with three samples per site. 100 g of each sample were collected and placed in sterile plastic bags. Samples were then maintained at +4°C until use.

Phytopathogenic bacteria isolation and identification

The serial dilution method was applied to isolate *Ralstonia solanacearum* from soil. One g of soil sample was added to 9 ml of sterile distilled water. Soil suspension was shaken in rotary shaker at 160 rpm until homogenization and suspensions were plated in Kelman's triphenyl tetrazolium chloride (TZC) agar medium [4] supplemented with one percent of yeast extract. Plates were incubated for 48h at 28°C. Bacterial colonies developing the typical irregular mucoid of *Ralstonia solanacearum* morphotype were subcultured on fresh TZC medium for further purification. Pure isolates were, subjected to phenotypic identification [5], biovar determination [6] and molecular characterization.

PCR amplification for species confirmation

Genomic DNA of isolates was extracted using DNeasy Blood & Tissue Kit (Qiagen, Cat. No. 69504) according to the manufacturer's instructions. A singleplex PCR with the *Ralstonia solanacearum* - specific primers 759/760 (5' GTC GCC GTC AAC TCA CTT TCC 3' and 5' GTC GCC GTC AGC AAT GCG GAA TCG 3') [7] was carried out in a 25 µl reaction volume with the following reagents: ~2 ng/µl of template DNA, 0.625 U of GoTaq Flexi, 4 pmoles of each primer, 1.5 mM MgCl₂, 0.2 mM dNTP (Promega) and 1X supplied amplification buffer PCR. The reaction was performed at the following steps: a denaturation step at 96°C for 5 min, 30 cycles at 94°C for 15 s, 29°C for 30 s and 72°C for 30 s, followed by a final extension at 72°C for 10 min. The expected 280-bp reference PCR products from the ribosomal DNA specific of *Ralstonia solanacearum* species were revealed by electrophoresis through 1.5% agarose gel in 50X TAE (Tris-acetate-EDTA) buffer stained with ethidium bromide and gels were imaged in GeneSnap (a Syngene Bioimaging System, Syngene) with UV transilluminator at a wavelength of 302 nm [7].

Isolation of antagonistic isolates

Fluorescent *Pseudomonas* spp. were isolated and counted using the King *et al.* technique [8]. This technique is based on the observation of pyoverdine under ultraviolet light at 254 nm, a fluorescent pigment which represents the majority of siderophores produced by these bacteria. For this, 1 g of each soil sample is weighed

and then serial dilutions (with 9 ml of PBS) were performed. Inoculation was done on the surface; inoculated plates were incubated at 28°C. All fluorescent colonies are counted and collected using a pre-sterilized “toothpick” and then subcultured into new plates containing King B medium.

Endophytic actinomycetes from leaves, stems and roots of healthy potato plant were isolated according to the method described by Fischer *et al.* [9].

Plant samples were washed in water and surface sterilized in ethanol 70% for 30 s. Then, they were treated with sodium hypochlorite (3%) for 3 min followed by three successive washings in sterile water. Each sample was divided into pieces of ca. 1 cm and pieces were aseptically transferred into petri dishes containing Starch Casein Agar medium (SCA) [10].

Actinomycete isolation from rhizospheric soil samples was performed by soil dilution plating technique on Starch Casein agar [11].

Selected colonies of actinomycetes (rough, chalky) were subcultured by incubating at 30°C and maintained on Waksman agar slants. Cultures were then kept at room temperature and 4°C or -20°C. Strains were subcultured to fresh media every two months.

***In vitro* antagonism assay**

Antagonistic activity of actinomycete and *Pseudomonas* isolates against the most pathogenic *Ralstonia solanacearum* strain R15 (pathogenicity assay not shown) was tested according to Almoneafy's method [12].

A culture of *Ralstonia solanacearum* was grown overnight in nutrient broth and then centrifuged. The cell pellet was suspended in sterilized saline solution (0.85% NaCl). Ten ml of *Ralstonia* suspension (10^8 CFU/ml) were mixed with 1 l of melted YPGA medium added with 5 ml of 2, 3, 5-triphenyl tetrazolium chloride (TZC), poured into sterile plates and allowed to solidify.

Thereafter, tested isolates were transferred to the surface of YPGA medium with sterile toothpicks and incubated at 28°C for four days for actinomycetes and two days for *Pseudomonas*. Four replicates were done under completely randomized design (CRD) for this assay. The inhibition of *Ralstonia* growth was revealed by the presence of clear halos without pinkish coloration around the tested isolates.

***In planta* assay**

Four isolates were selected for further test based on their *in vitro* activity against *Ralstonia solanacearum*. Glasshouse experimentation was used to test their ability to suppress potato bacterial wilt. Inocula were prepared by growing tested antagonistic isolates on SCA (actinomycetes) and on King B media (fluorescent *Pseudomonas*). Plates were then incubated at 30°C for two weeks for actinomycetes and two days for *Pseudomonas*.

The potato variety “Spunta” which is highly susceptible to *Ralstonia solanacearum*, was used in this assay. Cuttings were sterilized in sodium hypochlorite solution 2% for 2 min and washed thoroughly in sterile water. Samples were, then, soaked overnight in the bacterial suspension (10^8 CFU/ml) and cuttings immersed in sterile saline served as controls. Thereafter, potatoes inoculated with bacteria were planted into pots containing sterile soil mixed with peat and infected with the most pathogenic *Ralstonia solanacearum* strain (R15). The pots were placed in a greenhouse and maintained at 30°C and, 60% humidity in order to favor disease development. Furthermore, two control treatments were prepared: control one (C1) where cuttings were treated with pathogen only and control two (C2) where cuttings were planted without pathogen or tested antagonistic isolates. Every treatment was run in four replicates with five plants in each pot. All treatments were distributed in a randomized position to avoid any positional effect on plant growth.

The treated plants were monitored for disease development for 30 days and disease index was rated using the following scale: DI 0: no wilting; DI 1: 1-25% wilt symptom; DI 2: 26-50% wilt symptom; DI 3: 51-75% wilt symptom; DI 4: 76-100% wilt symptom or dead [13].

The disease incidence and biocontrol efficacy were estimated accordingly [14]:

Disease incidence (%) = [(Disease index x number of diseased plant in this index)/(Total number of plants investigated x the highest disease index)] x 100.

Biocontrol efficacy (%) = [(Disease incidence of control - Disease incidence of antagonist-treated group)/(Disease incidence of control)] x 100.

Plant growth parameters evaluation

After two months of treatment, the effect of antagonistic isolates on plant growth was assessed in terms of plant height, fresh and dry biomass weight. Dry biomass was obtained by drying plant samples at 60°C for three days. Moreover, growth promotion efficacy (GPE) was calculated to show the relative effect of antagonistic isolates on plant growth compared with the two control treatments by the following formula [12]:
Growth promotion efficacy (%) = [(Growth parameter in antagonist-treated group - Growth parameter in control group)/(Growth parameter in control group)] x 100.

Characterization of active bacterial strains

Three actinomycetes isolates that showed high antagonistic activity and enhanced plant growth in greenhouse were characterized morphologically following the directions given by the International Streptomyces Project (ISP) [15] and Bergey's Manual of Systematic Bacteriology [16]. Cultural characteristics of pure isolates in various media: ISP2, ISP4, ISP5, Starch Casein Agar were recorded after incubation for seven to 21 days at 30°C. Morphological observations were made using a light microscope [15].

Statistical analysis

The GenStat software was used for data analysis [17]. Mean comparisons were conducted using a Least Significant Difference (LSD) test ($P < 0.05$).

RESULTS

Isolation and characterization of Ralstonia solanacearum strains

Thirty strains were isolated from six soil rhizospheric samples under diseased potato plants based on morphological and cultural characteristics. Among these, 24 strains not showing fluorescent pigment production were subjected to biochemical characterization including biovar determination. The results given in Table 1 show that all isolates have catalase and oxidase enzymes. They use disaccharides including cellobiose, maltose and lactose but not sugar alcohols. On the basis of biochemical tests, strains are grouped into biovar two and PCR results show that the 759/760 primer set amplified a 280-bp fragment for all strains, supporting their identity as *Ralstonia solanacearum* (Figure 1).




Figure 1: DNA bands of *Ralstonia solanacearum* (280pb) in agarose gel after PCR amplification M: 1kb sizing ladder; R.s+: *Ralstonia solanacearum* reference strain (GMI1000), R.s-: negative control (no template)

Isolation and in vitro screening of antagonistic isolates

A total of 77 strains were obtained from two varieties of potato vegetal material and its rhizospheric soils (40 telluric actinomycetes, 25 endophytic actinomycetes and 12 fluorescent *Pseudomonas* spp.). Among 77 isolates screened, ten isolates P7, P10, S18, S19, S23, S25, E4, E7, E12 and E13 inhibited *Ralstonia solanacearum* strain development. Out of these, four isolates E7, S25, P7 and E13 gave a large zone inhibition with 40, 34, 26 and 23 mm diameter respectively; while the remaining isolates presented an inhibition zone from 9 to 20 mm of diameter (Figure 2).




Figure 2: Diameter of inhibition zone (mm) of ten antagonistic bacterial strains on *Ralstonia solanacearum* E: Endophytic actinomycetes; S: Telluric actinomycetes; P: Fluorescent *Pseudomonas*

In vivo evaluation of potential antagonists

Antagonistic bacterial strains showed significant differences in their ability to suppress potato bacterial wilt. Isolate E7 showed the lowest value of wilt incidence as well as the highest value of biocontrol efficacy against *Ralstonia solanacearum* with 20.08 and 72.04%, respectively. However, isolate P7 exhibited the highest disease incidence and the lowest value of biocontrol efficacy with 68.98 and 4.53%, respectively. Isolates E13 and S25 sustained significantly lower disease incidence and biocontrol efficacy compared with control (Table 2).

Indeed, significant differences among treatments regarding plant height and biomass were noted. The highest values were recorded for plants treated with the isolates E13 and S25 for all measured parameters compared to the control and other treatments (Table 3). The weights of fresh shoot biomass were 8.75 and 9.118 g, respectively with a corresponding GPE 68.75 and 75.85%. Those of dry shoot biomass were, however, 3.64 and 3.15 g with a GPE 115.42 and 86.7%, respectively. Otherwise, plant heights were 60.66 and 60.75 cm for treatments with E13 and S25. Their GPE were 44.63 and 44.84%, respectively.

Cultural and morphological characterization of selected isolates

The three isolates E7, E13 and S25 grew on a range of agar media showing morphology typical of Streptomycetes. The color of the substrate mycelium and aerial spore mass varied (brown and white for the isolates E7 and S25, red for the isolate E13). The isolate E13 produced diffusible pigment on SCA agar media. Melanin was produced on peptone-yeast extract agar (ISP6) and ISP7 agar.

Aerial hyphae of all isolates differentiated into long spiral chains of cylindrical spores. The vegetative hyphae of all isolates were branched but not fragmented. Verticils were not detected.

DISCUSSION

From two infected potato fields in Antsirabe II district in the Vakinankaratra region of Madagascar, 30 phytopathogenic strains of *Ralstonia solanacearum* were isolated from six soil samples. Polymerase chain reaction (PCR) amplification with specific primers confirmed that the isolates were *Ralstonia solanacearum* strains and biochemical test showed that they belong to the biovar two. Thus, our result is different from those reported by some authors [18, 19] in a survey of potato plants for bacterial wilt in Madagascar, where *Ralstonia solanacearum* was found to belong to the biovar one. To our knowledge, this is the first record of *Ralstonia solanacearum* strain, biovar two in Madagascar. However, further investigation is necessary for origin confirmation and phylotype classification.

An Additional, 77 strains for biocontrol test were isolated from healthy plants and their rhizospheric soils. Among these, antagonism assays showed that ten isolates displayed remarkable antagonistic activity against *Ralstonia solanacearum* and the isolates E7, E13, S25 and P7 exhibited the highest inhibition effect. Some previous studies reported that actinomycetes and fluorescent *Pseudomonas* spp. showed antagonistic activity against phytopathogenic bacteria and fungi [20, 21, 22]. This antagonistic mechanism may be due to antibiosis or production of siderophores or both [23]. Moreover, endophytic actinomycetes were found to produce at least three kinds of antagonistic substances in plant tissues including antibiotics, enzymes and siderophores [24]. The inhibition of pathogen bacteria development could be due to the emission of volatile secondary metabolites secreted by actinomycetes.

In greenhouse experimentation, the application of two endophytic actinomycete strains and one telluric actinomycete on potato plants induced good reduction of disease severity of potato bacterial wilt. This reduction was significantly higher than observed with the *Pseudomonas* isolate. Furthermore, plants treated with these three strains showed high values of plant height and biomass (fresh and dry weight) compared with the control treatment. Previous work has demonstrated that root-associated bacteria are promising tools to control soil-borne pathogens and to enhance plant growth [25]. This is in agreement with our study which shows that E7, E13 (endophytic actinomycetes isolated from potato roots) are active against *Ralstonia solanacearum* and significantly improve plant growth.

Plant growth promotion may be due to the production of plant growth promoters by efficient strains. Several species of *Streptomyces* secrete indole-three-acetic acid (IAA) when fed with L-tryptophan and improve plant growth [26]. It has been also demonstrated that some endophytic actinomycetes produce plant growth promoters with auxin-like activity; these compounds were shown to accelerate the formation of plant adventitious roots [27].

One isolate of *Pseudomonas* (P7) showed high antagonism *in vitro* but resulted in the lowest disease suppression in planta. A previous study demonstrated that fluorescent *Pseudomonas* may succeed as a biocontrol agent *in vitro* or under controlled conditions but fail under pot experiments and field conditions [28]. This could be explained by the fact that the development of the biocontrol agents is influenced by environmental factors such as unfavorable conditions for growth and competition with other microorganisms.

Cultural and morphological characterization of the three active isolates using a light microscopy showed that they belong to the genus *Streptomyces*. Branching filaments, abundant aerial mycelia, and long chains of small rectiflexible spores are visible, all of which are characteristic of *Streptomyces* sp. [29]. These isolates showed opaque, rough, nonspreading colonies which adhere to agar medium as described in Bergey's Manual of Determinative Bacteriology [16, 30]. Morphological examination indicated that the three isolates belong to the genus *Streptomyces*.

In conclusion, this study reports for the first time the antagonistic activity of actinomycetes and *Pseudomonas* strains against *Ralstonia solanacearum*, the causal agent potato bacterial wilt in Madagascar. Therefore, further work will be needed to investigate the mode of action of the strains that showed high antagonistic activity and high performance to enhance plant growth (E7, E13 and S25) and to confirm their effectiveness under natural conditions.

ACKNOWLEDGEMENT

This work was funded by PARRUR and SCAC of France Embassy. We would like to thank the whole *Ralstonia* project team: FIFAMANOR, CNRE, CIRAD La Réunion, Biochemistry and Entomology Departments of the University of Antananarivo. We are also grateful to Dr Leah Blasiak and Dr Liva Rakotondraibe for helping to revise the paper.

Table 1: Identification of isolates based on biochemical characterization and biovar determination

Isolate	Biochemical test		Biovar determination					
	Character		Disaccharides and sugar alcohols					
	Catalase	Oxydase	Mal	Lac	Cel	Dul	Man	Sor
R1-R2-R3-R4-R5-R6-R7- R8- R9- R12-R13-R14 R15-R17-R18-R20-R21- R22-R23-R25-R26-R27- R28-R30	+	+	+	+	+	-	-	-

Mal: Maltose; **Lac:** Lactose; **Cel:** Cellobiose; **Duc:** Dulcitol; **Man:** Mannitol; **Sor:** Sorbitol; **R:** *Ralstonia* strains
+ indicates presence and – indicates absence of the character

Table 2: Disease incidence and biocontrol efficacy of treatments with four isolates E7, E13, S25 and P7 against *Ralstonia solanacearum*

Treatment	Disease incidence (%)	Biocontrol efficacy (%)
E7+R.s	20.08 ± 1.46 ^c	72.04 ± 1.14 ^a
E13+R.s	50.37 ± 1.62 ^b	30.60 ± 1.56 ^c
S25+R.s	47.26 ± 0.99 ^b	34.41 ± 2.13 ^b
P7+R.s	68.98 ± 1.27 ^a	4.53 ± 0.91 ^d
R.s	71.06 ± 0.91 ^a	-

Mean followed by the same letter within a column are not significantly different as determined by the LSD test ($p < 0.05$)

Table 3: Effect of application of antagonistic bacterial isolates and *Ralstonia solanacearum* on the height and weight of fresh and dried potato plants

Treatment	Fresh shoot biomass (g)		Dry shoot biomass (g)		Plant height (cm)	
	Mean	GPE (%)	Mean	GPE (%)	Mean	GPE (%)
E7+R.s	5.91 ± 0.25 ^d	13.87	2.05 ± 0.06 ^c	21.33	52.11 ± 1.57 ^b	24.24
E13+R.s	8.75 ± 0.48 ^a	68.75	3.64 ± 0.43 ^a	115.42	60.66 ± 1.27 ^a	44.63
S25+R.s	9.11 ± 0.13 ^a	75.85	3.15 ± 0.09 ^b	86.17	60.75 ± 1.28 ^a	44.84
P7+R.s	3.94 ± 0.28 ^f	-	1.84 ± 0.13 ^c	8.92	48.66 ± 0.96 ^c	16.02
R.s	5.18 ± 0.36 ^e	-	1.69 ± 0.09 ^d	-	41.94 ± 1.14 ^e	-
Saline solution	6.77 ± 0.19 ^c	-	2.97 ± 0.09 ^b	-	46.01 ± 0.77 ^d	-

Mean followed by the same letter within a column are not significantly different as determined by the LSD test ($p < 0.05$); Data represent means of the experiment with four replications each.

GPE: Growth Promotion Efficacy

R.s: *Ralstonia solanacearum*

REFERENCES

1. **Hayward AC** *Ralstonia solanacearum*. **In:** Encyclopedia of Microbiology. J. Lederberg, ed. Academic Press, San Diego, CA. 2000.
2. **Crawford DL, Lynch JM and MA Ousley** Isolation and characterization of actinomycetes antagonists of a fungal root pathogen. *Appl. Environ. Microbiol.* 1993; **59** (11): 3899-3905.
3. **Curl EA and B Truelove** The rhizosphere. Advanced Series in Agriculture Sciences, 15. Springer-Verlag Berlin Heidelberg. 1986; 288 pp.
4. **Kelman A** The relationship of pathogenicity in *Pseudomonas solanacearum* to colony appearance on tetrazolium medium. *Phytopathol.* 1954; **44**:693-695.
5. **Zubeda C and R Hamid** Isolation and characterization of *Ralstonia solanacearum* from infected tomato plants of soanskesar valley of punjab. *Pak. J. Bot.* 2011; **43**(6): 2979-2985.
6. **Denny TP** Plant pathogenic *Ralstonia* species. **In:** Bacteria SS and Gnanamanickam (Ed). Plant-Associated, Springer Publishing, Dordrecht, The Netherlands. 2006: 573-644.
7. **Opina N, Tavner F, Hollway G, Wang JF, Li TH, Maghirang R, Fegan M, Hayward AC, Krishnapillai V, Hong WF, Holloway BW and JN Timmis A** novel method for development of species and strain specific DNA probes and PCR primers for identifying *Burkholderia solanacearum* (formely *Pseudomonas solanacearum*). *Asi. Pac. Mol. Biol. Biotechnol.* 1997; **5**: 19-30.
8. **King EO, Ward MKM and DE Raney** Two simple media for the demonstration of pyocyanin and fluorescein. *J. Lab. Clin. Med.* 1954; **44**: 301-307.
9. **Fisher PJ, Petrini O and HM Lappin-Scott** The distribution of some fungal and bacteria endophytes in maize (*Zea mays* L.). *New. Phytol.* 1992; **122**: 299-305.
10. **Kuster E and ST Williams** Selection of media for isolation of Streptomycetes. *Nat.* 1964; **202**: 928- 929.
11. **Oskay M, Usame A and C Azeri** Antibacterial activity of some actinomycetes isolated from farming soils of Turkey. *Afr. J. Biotechnol.* 2004; **3**: 441-446.
12. **Almoneafy AA, Xie GL, Tian WX, Xu LH, Zhang GQ and M Ibrahim** Characterization and evaluation of *Bacillus* isolates for their potential plant growth and biocontrol activities against tomato bacterial wilt. *Afr. J. Biotechnol.* 2012; **11**(28): 7193-7201.

13. **Park EJ, Lee SD, Chung EJ, Lee MH, Um HY, Murugaiyan S, Moon BJ and SW Lee** Microtom- a Model Plant System to Study Bacterial Wilt by *Ralstonia solanacearum*. *Plant Pathol. J.* 2007; **23(4)**: 239-244.
14. **Xue QY, Chen Y, Li SM, Chen LF, Ding GC, Guo DW and JH Guo** Evaluation of the Strains of *Acinetobacter* and *Enterobacter* as Potential Biocontrol Agents against *Ralstonia* Wilt of Tomato. *Biol. Control.* 2009; **48(3)**: 252-258.
15. **Shirling EB and D Gottlieb** Methods of characterization of *Streptomyces* species. *Int. J. Sys. Bacteriol.* 1996; **16**: 313-349.
16. **Locci R** *Streptomyces* and related Genera. **In:** Bergey's Manual of Systematic Bacteriology. Williams & Wilkins Company, Baltimore. 1989; **4**: 2451-2508.
17. **Buyse W, Stern R and R Coe** Genstat Discovery Edition for everyday use. ICRAF editor. Nairobi, Kenya. 2004.
18. **Prunier JP** Différenciation en types biochimiques de souches de *Pseudomonas solanacearum* (Smith) Smith. *Phitiat. Phytophar.* 1967; **16**: 199-204.
19. **Lallmahomed GM, Rakotobe R and Rakotondramanana** Biovars and races of *Pseudomonas solanacearum* in Madagascar - a preliminary study. *FAO Plant. Prot. Bill.* 1988; **36(2)**: 54-59.
20. **El Albyad MS, el Sayed MA and AR el Shanshoury** Effect of culture conditions on the antimicrobial activities of UV-mutants of *Streptomyces corchorusii* and *Streptomyces spiroverticillatus* against bean and banana wilt pathogens. *Microbiol. Res.* 1996; **151**: 201-211.
21. **Guo JH, Guo YH, Zhang LX, Qi HY and ZD Fang** Screening for biocontrol agents against cayenne pepper bacterial wilt. *Chi. J. Biol. Control.* 2001; **17**: 101-106.
22. **Götz M, Gomes NCM, Dratwinski A, Costa R, Berg G, Peixoto R, Mendonça-Hagler L and K Smalla** Survival of gfp-tagged antagonistic bacteria in the rhizosphere of tomato plants and their effects on the indigenous bacterial community. *FEMS Microbiol. Ecol.* 2006; **56**: 207-218.
23. **Lemessa F and W Zeller** Screening Rhizobacteria for Biological Control of *Ralstonia solanacearum* in Ethiopia. *Biol. Control.* 2007; **42(3)**: 336-344.
24. **Trejo-Estrada SR, Rivas SI and D L Crawford** *In vivo* and *In vivo* Antagonism of *Streptomyces violaceusniger* YCED9 against Fungal Pathogens of Turfgrass. *World. J. Microbiol. Biotechnol.* 1998; **14**: 865-827.
25. **Weller DM** 1988. Biological-Control of Soilborne Plant-Pathogens in the Rhizosphere with Bacteria. *Ann. Rev. Phytopathol.* 1988; **26**: 379-407.

26. **Manulis S, Shafrir H, Epetein E, Lichter A and I Barash** Biosynthesis of indole-3-acetic acid via indole-3-acetamide pathway in *Streptomyces* spp. *Microbiol.* 1994; **140**: 1045-1050.
27. **Igarashi Y, T Iida R, Yoshida and T Furumai** Pteridic acids A and B, novel plant growth promoters with auxin-like activity from *Streptomyces hygroscopicus* TP-A0451. *J. Antibiot.* 2002; **55**: 764 –767.
28. **Ran LX, Liu CY, Wu GJ, Van Loon LC and PAHM Bakker** Suppression of bacterial wilt in *Eucalyptus urophyllaby* fluorescent *Pseudomonas* spp. in China. *Biolog. Control.* 2005; **32**: 111-120.
29. **Lechevalier HA** The Actinomycetes III, A Practical Guide to Generic Identification of Actinomycetes. **In:** Bergey's Manual of Systematic Bacteriology. Williams & Wilkins Company, Baltimore. 1989; **4**: 2344-2347.
30. **Anderson AS and MHE Wellington** The taxonomy of *Streptomyces* and related genera. *Int. J. Syst. Evol. Microbiol.* 2001; **51**: 797-814.