

# Morphology of the larvae, male genitalia and DNA sequences of *Anopheles (Kerteszia) pholidotus* (Diptera: Culicidae) from Colombia

Jesús Eduardo Escobar<sup>1,2/+</sup>, Ranulfo González<sup>3</sup>, Martha L Quiñones<sup>1</sup>,  
Richard C Wilkerson<sup>4</sup>, Fredy Ruiz<sup>4</sup>, Bruce A Harrison<sup>5</sup>

<sup>1</sup>Universidad Nacional de Colombia, Bogotá, Cundinamarca, Colombia <sup>2</sup>Universidad de la Salle, Bogotá, Cundinamarca, Colombia  
<sup>3</sup>Faculty of Natural and Exact Sciences, Universidad del Valle, Valle, Colombia <sup>4</sup>Walter Reed Biosystematics Unit, Smithsonian Institution, Museum Support Center, Suitland, MD, USA <sup>5</sup>College of Health and Human Sciences, Western Carolina University, Clemmons, NC, USA

Since 1984, *Anopheles (Kerteszia) lepidotus* has been considered a mosquito species that is involved in the transmission of malaria in Colombia, after having been incriminated as such with epidemiological evidence from a malaria outbreak in Cunday-Villarrica, Tolima. Subsequent morphological analyses of females captured in the same place and at the time of the outbreak showed that the species responsible for the transmission was not *An. lepidotus*, but rather *Anopheles pholidotus*. However, the associated morphological stages and DNA sequences of *An. pholidotus* from the foci of Cunday-Villarrica had not been analysed. Using samples that were caught recently from the outbreak region, the purpose of this study was to provide updated and additional information by analysing the morphology of female mosquitoes, the genitalia of male mosquitoes and fourth instar larvae of *An. pholidotus*, which was confirmed with DNA sequences of cytochrome oxidase I and rDNA internal transcribed spacer. A total of 1,596 adult females were collected in addition to 37 larval collections in bromeliads. Furthermore, 141 adult females, which were captured from the same area in the years 1981-1982, were analysed morphologically. Ninety-five DNA sequences were analysed for this study. Morphological and molecular analyses showed that the species present in this region corresponds to *An. pholidotus*. Given the absence of *An. lepidotus*, even in recent years, we consider that the species of mosquitoes that was previously incriminated as the malaria vector during the outbreak was indeed *An. pholidotus*, thus ending the controversy.

Key words: *Kerteszia* - *Anopheles pholidotus* - male genitalia - DNA sequences

Mosquito species of the subgenus *Kerteszia* are exclusively located in the Neotropical areas of Central and South America. Except for Chile and Uruguay, their distribution ranges from the south of Mexico to southern Brazil. This subgenus is also present in the Caribbean, Trinidad and Tobago (de Carvalho-Pinto & Lourenço-de-Oliveira 2004) and both the Atlantic and Pacific coastal areas (Marrelli et al. 2007), with some species present at altitudes of more than 1,000 m above sea level (a.s.l.) (Cova-García 1961, Harbach & Navarro 1996).

*Kerteszia* subgenus consists of species that have been incriminated as malaria vectors and are related to the emergence of malaria in thermal floors at altitudes of over 1,500 m a.s.l. Moreover, in the malaria-heavy coastal regions of South America and in the eastern slopes of the Venezuelan Andes, members of this subgenus seem to be the most important species involved in the transmission of malaria (Benítez et al. 2004, Montoya-Lerma et al. 2011). Similarly, the species of *Kerteszia* have been incriminated in the transmission of “malaria by bromeliads”, which is typical of protected areas because the

epiphytic bromeliads are the only breeding sites. This is why these plants have been associated with the occurrence of the disease (Ueno et al. 2007).

Five of the 12 species described for this subgenus (Zavortink 1973, Collucci & Sallum 2003) have been incriminated as malaria vectors: *Anopheles bellator* (Forattini et al. 1999), *Anopheles cruzii* (de Carvalho-Pinto & Lourenço-de-Oliveira 2004), *Anopheles homunculus* (Rubio-Palis & Zimmerman 1997), *Anopheles neivai* (Carvajal et al. 1989, Gutierrez et al. 2008) and *Anopheles lepidotus* (Quiñones et al. 1984, Montoya-Lerma et al. 2011).

In Colombia, seven species of the subgenus have been reported, among which *An. neivai* and *An. lepidotus* are considered malaria vectors, while *Anopheles boliviensis* is considered a seasonal vector only (Quiñones et al. 1984, Montoya et al. 1994, Olano et al. 2001).

Since 1984, *An. lepidotus* has been considered, using epidemiological evidence (namely, mosquito density correlated with malaria cases; 98% of the *Anopheles* species found in the malaria focal area corresponded to this species), to be one of the species responsible for malaria transmission in Colombia, particularly in the focal area of Cunday-Villarrica, Department of Tolima. It has been suggested that *An. lepidotus* is part of a complex of species in which members of this complex are difficult to be differentiated morphologically. Specimens of *Kerteszia* from Ecuador and Venezuela have been analysed using molecular tools, thereby providing evidence for a complex of species: *An. lepidotus* could exist with *Anopheles pholidotus* and *An. boliviensis* in a complex of at least

doi: 10.1590/0074-0276130596

Financial support: COLCIENCIAS (110145921478)

+ Corresponding author: jeescobar@unisalle.edu.co

Received 26 December 2013

Accepted 8 April 2014

four or five different species (R Wilkerson, unpublished observations, Sallum et al. 2002).

González and Carrejo (2009), using observations of specimens of what was supposedly *An. lepidotus*, noted that the mosquito is very similar to *An. pholidotus* and both species were called *An. boliviensis* before Zavortink's (1973) descriptions. As a result, the holotype and paratype described by Komp (1937) as *An. boliviensis* is actually *An. lepidotus*. They also affirm that of the 204 Colombian specimens deposited at the University of California, Los Angeles and the National Museum of Natural History (NMNH) at the Smithsonian Institution collections and predetermined to be *An. boliviensis* as analysed by Zavortink (1973), 85% corresponded to *An. lepidotus*. Similarly, González and Carrejo (2009), based on the analysis of male genitalia and larvae of four specimens from Cunday-Villarrica, suggested the possibility that *An. pholidotus* occurred in this region. The importance of the above conclusion is that in Colombia, the region of Cunday-Villarrica is considered to be a focal area of malaria, where *An. lepidotus* had been incriminated as the possible vector for more than 25 years (Quiñones et al. 1984).

Recently, Harrison et al. (2012) resolved the problem of separating *An. lepidotus* from *An. pholidotus* females by re-describing both species and preparing a taxonomic key. These methods were used to differentiate females, the IV instar larvae, pupae and male genitalia of *An. lepidotus* from other species of *Kerteszia*. They examined specimens of *An. lepidotus* from Colombia, Ecuador and Peru and female specimens of *An. pholidotus* from Bolivia, Colombia, Costa Rica, Ecuador and Venezuela. They agree with the preliminary results of Escobar et al. (2010), who stated that the primary vector in Tolima is *An. pholidotus* and not *An. lepidotus*, as had been reported since 1984 (Quiñones et al. 1984). Nevertheless, Harrison et al. (2012) only analysed 33 females of *An. pholidotus* captured between 1981-1983 in Tolima. Therefore, this conclusion was pending the analysis of the morphology of associated stages, the male genitalia and the inclusion of DNA sequences from samples of *An. pholidotus* that were recently obtained from the study area.

Since the controversy surrounding the identity of the vector in Tolima began more than 30 years ago, the aim of this study was to provide updated and additional information that was not presented by Harrison et al. (2012). This information includes the morphology of larvae, the male genitalia of *An. pholidotus* and their association with the DNA sequences of specimens that were recently collected in Colombia.

In this study, we analysed the morphological characteristics of 1,596 recently captured females, the chaetotaxy of IV instar larvae and the male genitalia of specimens of *An. pholidotus* from three localities in Tolima. Molecular analysis for gen cytochrome oxidase I (*COI*), internal transcribed spacer (ITS2) and sequencing was performed to verify the results obtained from the morphological information.

This study provides definitive morphological and molecular information supporting the conclusion that the malaria vector in Tolima is *An. pholidotus* and that it is actually the only malaria vector present in the area.

## MATERIALS AND METHODS

**Study site** - The malaria focal area of Cunday-Villarrica is composed of the municipalities of Cunday, Villarrica, Dolores, Prado and Purificación (03°57'N 074°36'W) (Fig. 1), which are situated on the western slopes of the eastern mountain chain and have an average temperature of 24°C and an annual rainfall of between 2,000-4,000 mm. This area is approximately 100 Km<sup>2</sup> and is characterised by humid forests that have an abundance of epiphytes. This area has had stable malaria transmission for more than 25 years. Moreover, malaria cases reported in Tolima have been almost exclusively from that focal area.

**Sampling strategy and adult mosquitoes** - Collections of biological material were made between February 2009-June 2012. Adult females were captured using human landing collections inside and around houses and in the forest during periods of peak biting at various locations in the municipalities in this study. Captured females were individualised in vials and brought to the entomology laboratory of the National University of Colombia for preservation and taxonomic determination. Additionally, adult females of *Kerteszia*, which were collected between 1981-1983 in the same study area and belonged to the collection of the entomology laboratory of the Museum of Entomology, University of Valle, were analysed morphologically.

**Entomological series** - Mosquitoes in immature stages (larva and pupa) were captured directly from water at the base of bromeliads leaves by drawing out their




Fig. 1: location of the study area, foci of Cunday-Villarrica, Department of Tolima, Colombia. Map realised by Geógrafos del Valle.

content with a larger pipette. Larvae of *Anopheles* were collected from 41 trees with 78 bromeliads and brought to the laboratory of the National University of Colombia for rearing to the adult stage. IV instar larvae were individualised and larval and pupal skins were obtained to achieve 37 full entomological series. Male genitalia and larval and pupal skins were mounted.

**Morphological analysis** - Diagnostic characters of females, IV instar larvae and male genitalia were studied based on the descriptions and morphological keys of Zavortink (1973) and Harrison et al. (2012).

**Molecular methods** - Forty-four specimens were processed as belonging to *An. (Kerteszia)* sp. and their DNA was extracted using the DNeasy Blood & Tissue Kit (QIAgen®, USA). The rDNA ITS2 region was amplified using the primers of Collins and Paskewitz (1996) and the polymerase chain reaction (PCR) conditions described by Linton et al. (2001). A portion (710 bp) of the barcoding region of *COI* (mDNA) was amplified using the primers designed by Folmer et al. (1994) and the PCR conditions described by Ruiz et al. (2010). The products were visualised on 1% agarose gel containing 0.5 mg/mL of ethidium bromide. The PCR product was purified using ExoSAP-IT® (USB Corporation, USA).

Sequencing reactions were carried out in both directions using the Big Dye terminator Kit® (PE Applied Biosystems, England) on an ABI 3730 automated sequencer (PE Applied Biosystems). The sequences were edited with Sequencher™ 4.10.1 (Gene Codes Corporation, USA) and aligned manually and translated to amino acids in MacClade v.4.06 (Maddison & Maddison 2003). Sequence similarities were compared with those available in GenBank using Basic Local Alignment Search Tool (BLAST) (ncbi.nlm.nih.gov/genbank/) and sequence statistics were calculated in MEGA v.5 (Tamura et al. 2011).

**Ethics** - This study was approved by the Ethical Committee of the Faculty of Medicine of the National University of Colombia, according to consecutive number E-31, on 26 June 2008.

## RESULTS

Table shows the sampling locations in the study area and the sampling dates. A total of 1,737 anopheline females were analysed morphologically. Of these females, 1,596 were collected landing on humans in the period between 2009-2012 and 141 females were sampled between 1981-1983.

The first morphological analysis was conducted using the dichotomous keys of Zavortink (1973). As a result of this analysis, 76% (1,218) of the collected specimens corresponded to either *An. lepidotus* or *An. pholidotus* and 24% (378) were in agreement with an identification of *An. boliviensis*. According to Zavortink's key (1973), the only difference existing between *An. lepidotus* and *An. pholidotus* is restricted to the size of the scales on the female abdomen. As a result, it is difficult to determine which of these two species is present in the study region using this morphological key. Considering this issue, a second morphological analysis was made using

the dichotomous key proposed by Harrison et al. (2012). This key can separate *An. lepidotus* females from *An. pholidotus* females according the following diagnostic features: (i) females with white scales on the proboscis, pedicel and palpomere I; hindtarsomeres I and 2 without apical pale band (from dorsal view) are classified as *An. lepidotus* and (ii) females without white scales on the proboscis, pedicel and palpomere I; hindtarsomeres I and 2 with narrow apical pale band (from dorsal view) are classified as *An. pholidotus*.

All females captured in Tolima showed morphological characteristics that were similar to those assigned to *An. pholidotus*. Additionally, the females of *An. pholidotus* exhibited numerous scales with variable widths on tergites and sternites II-VII, hindtarsal segment 2 with white bands in the apical pale band between 0.1-0.2 as the total length of the tarsomere, mesanepimeron with large curved patch of scales extending from higher bristles to below the middle of the segment, scales in the proximal tergites and scales of the distal tergites forming transverse apical bands (Fig. 2).

**IV instar larvae** - Twenty-eight micro-preparations of IV instar larvae were analysed. The larvae showed a characteristic coloration pattern in the dorsal area of the thorax and abdominal segments I and V (Fig. 3). The morphological characteristics of this instar coincided with those described by Zavortink (1973) for *An. pholidotus*. Setae 5-7-C simple, not plumose, moderately long, characteristics of the *Kerteszia* subgenus; setae 6-VI moderately long, always different to 6-III-V, setae 1-III-VII small palmate with pointed leaflets, pecten teeth of similar size all long, with marginal spinules extended to the apex, setae 1-I not palmate, setae 4-C always less developed than 2-C, setae 1-VII filiform, not palmate.

According to Zavortink (1973), two morphological characteristics help differentiate *An. lepidotus* from *An. pholidotus* in this larval stage: *An. pholidotus* shows a setae 3-C that is moderately developed and approximately 1/2 the length of 2-C, while *An. lepidotus* shows a setae 3-C that is short and thick, fusiform and less than 1/2 the length of 2-C. The 28 specimens analysed in this study presented an average length of  $0.19 \pm 0.01$  mm for the setae 2-C and of  $0.11 \pm 0.004$  mm for the setae 3-C, showing that in all cases the length of the setae 3-C is more than half the length of the setae 2-C (Fig. 4A), which corresponds to *An. pholidotus*. The clypeal index was  $0.98 \pm 0.15$ . The second discriminatory character is the setae 11-P, which is well developed for *An. pholidotus*. A very short 11-P would correspond to *An. lepidotus*. All larvae tested showed the setae 11-P to be well developed (length =  $0.32 \pm 0.03$  mm), with an average of 66.7% of the length of the 12-P setae ( $0.48 \pm 0.03$  mm) (Fig. 4B).

Some diagnostic characteristics described by Harrison et al. (2012) showed that for *An. lepidotus*, the seta 3-C is very thick, short and usually sharply pointed; however, seta 11-P is always very short and setae VI is stout and long. As mentioned above, the specimens analysed in this investigation showed characteristics that do not match those described for *An. lepidotus*.



TABLE  
Sampling localities of Cunday-Villarrica focus in the Department of Tolima, Colombia

Municipality	Coordinates	Locality	Sampling period
Cunday	04°47'59.1"N 74°34' 50.21"W	San Francisco	February 2009
Icononzo	04°51'02.1"N 74°09'003"W	Valencia Cuatimbal Alto Icononzo	November 1981 October 1982 October 1982
Villarrica	03°52'03.4"N 74°39'09.8"W	Puerto Lleras Totumal Guanacas Rio Lindo Rio Lindo Rio Lindo Rio Lindo	February 1981-December 1982 June 1981-February 1983 June 1982-February 1983 February-November 2009 August 2010 August-September 2011 April-July 2012
Purificación	03°51'58.11"N 74°55'59.83"W	Villa Esperanza	May 2009
Prado	03°45'04.15"N 74°49'55.9"W	El Cruce	May 2009



Fig. 2: lateral view of female of *Anopheles pholidotus*. Right arrow shows mesepimeron with one large and curved patch of scales that extends ventrally from upper setae. Left arrow shows palpomere 1 without scales.



Fig. 3: general view that shows the dorsal coloration pattern of larvae IV instar of *Anopheles pholidotus*.

**Male genitalia** - Twenty-five micro-preparations of male genitalia from a series were analysed. The morphological characteristics for this structure coincided with those described by Zavortink (1973) and González and Carrejo (2009) for *An. pholidotus*, but not for *An. lepidotus*. The specific characteristics were gonocoxite with long and sinusoid parabasal setae, always greater than 1/3 the total length of gonocoxite; one internal seta flattened at the apex and two accessory ones located apically, an aedeagus without leaflets (Fig. 5A), a lateral expansion of the ventral lobe of claspette, moderately spiculate except laterally and rounded (Fig. 5B, C). The 25 male genitalia analysed in this study showed a lateral expansion of the ventral lobe of claspette corresponding

to the typical shape of *An. pholidotus* described by Zavortink (1973) and González and Carrejo (2009).

**DNA sequences** - Ninety-five specimens were sequenced with the molecular markers *COI* and *ITS2*. Only one haplotype was observed for each marker, confirming the lack of an intraspecific variation in all of the individuals analysed.

The *COI* barcode region (658 bp) consisted of a single open reading frame, discarding the presence of pseudogenes. Its nucleotide frequencies (%) were A: 27.7%, T: 40.9%, G: 15.5% and C: 16% (Fig. 6). The *ITS2* fragment showed no length variability (546 bp) and its nucleotide frequencies were A: 26.4%, T: 19.4%, G: 26.7% and C: 27.5% (Fig. 7).




Fig. 4: larvae IV instar of *Anopheles pholidotus* (400X). A: head. Clipeal setae (2-C and 3-C); B: prothorax. Setae 11-P.



Fig. 5: male genitalia of *Anopheles pholidotus*. A: general view (200X) (ae: aedeagus; gc: gonocoxite; sa: accessory setae; si: internal setae; sp: parabasal setae); B: view of lateral expansion of ventral lobe of claspette (400X); C: ventral lobe of claspette expanded showing the characteristic form of *An. pholidotus* (1,000X).

The sequences of ITS2 were submitted to BLAST, where high significant homology (100%) was found with the sequences of *An. pholidotus* (from Táchira, Venezuela) (accessions JN967769.1, JN967768.1) published by Harrison et al. (2012).

*COI* sequences, compared with sequences in the Barcode of Life Data Systems (Bold Systems) (boldsystems.org), showed 100% similarity with *An. pholidotus*.

#### DISCUSSION

Until the study published by Harrison et al. (2012), the only morphological feature that had been proposed to differentiate females of *An. pholidotus* from *An. lepi-*

*dotus* were, according to Zavortink (1973), the size of the scales on proximal tergites. Those were described as moderately wide or broad for *An. lepidotus* and predominantly narrow to moderately wide for *An. pholidotus*, with those differences being rather subjective. This taxonomic characteristic is vague and difficult to interpret, so it would be inaccurate to separate female adults of these two species based only on these characteristics. Therefore, it was essential to analyse other morphological structures and associated stages, such as male genitalia and to use the chaetotaxy of IV instar larvae. Even so, genitalia analysis presents a “weak spot” or a “bottle neck” effect in the separation of these two species be-



GGTCAACAAA TCATAAGAT ATTGGTACTT TATATTTTAT TTTTGGAGCT TGAGCTGGAA 60  
 TAGTTGGAAC TTCTTTAAGA ATTTTAGTTC GAGTCGAATT AGGTCATCCT GGTGCTTTTA 120  
 TTGGAGATGA CCAAAATTAT AATGTTATTG TAACTGCTCA TGCTTTTATC ATAATTTTTT 180  
 TTATAGTAAT ACCTATATA ATTGGAGGAT TTGGAACATG ATTAGTTCCT TTAATCTTG 240  
 GAGCTCCTGA TATAGCTTTC CCTCGAATAA ATAATATAAG ATTTTGAATA CTTCCTCCTT 300  
 CTTAACTCT TTTAATTCA AGAAGTATAG TAGAAAATGG AGCTGGAACA GGTGGAACG 360  
 TTTATCTCC TCTATCATCT GGAATTGCC ACGCCGAGC TTCAGTAGAT TTAGCTATTT 420  
 TTTCTTTACA TTTAGCAGGT ATTTCTTCTA TTTTAGGAGC AGTAAATTTT ATTACCACAG 480  
 TAATTAATAT ACGATCTCCA GGAATTACTT TAGATCGAAT ACCATTATTT GTTGATCAG 540  
 TTGTAATTAC TGCAATTTTA TTACTTTTAT CTTTACCACT TTTAGCTGGA GCTATTACTA 600  
 TACTTTTAC TGACCGAAT CTTAATACTT CTTTTTTTGA CCCCGCTGGA GGAGGAGACC 660  
 CTATTTTATA CCAACATTTA TTTTGATT TTGGTCACCC TGAAGTTT 710

Fig. 6: cytochrome oxidase I sequence of *Anopheles pholidotus* from Villarrica, Department of Tolima, Colombia using the Folmer et al. (1994) primer (n = 10, 710 bp). Only one haplotype was found. Primers sequences are underlined and in bold.

ATCACTCGGC TCGTGGATCG ATGAAGACCG CAGCTAAATG CGCGTCAGAA TGTGAACATGC 60  
 AGGACACATG AACACCGATA CGTTGAACGC ATATTGCGCA TCGCACGACA CAGTCGATG 120  
 TACACATTTT TGAGTGOCCA TCCTCACCGC ATAGCCAATC ATCGGGGAAC GCGCCATGCG 180  
 CTTCCCGATG CATTATGATG CGTTGCCCG CCCCGCTGG TCAATCATG AAAGACTGTG 240  
 TGCGTAGGAG GGGCCGACCG GAGCGTCCG TTTGCGGAGT GTCTTCCGTC ACGGCTGAGC 300  
 CACCTTACGA CTCTACCAA CACGAGATAC GATGGTGAG AACAAATCCA TATCGAACGA 360  
 AACTACGCGT TCGGTGGCTG GCGAGTGCTG TTGACGATAG AGAGTTAGAG GAGAAGACAC 420  
 AGCAGGCAGA GAGACCGCG ATATCGTGG AGAGCGCGAA CGCACCGGCC GCGCTTACCA 480  
 CAACATGACA CACGAATCT ATAGTGGGCC TCAATAATG TGTGACTACC CCCTAAATT 540  
ATGCAT 546

Fig. 7: internal transcribed spacer sequence of *Anopheles pholidotus* from Villarrica, Department of Tolima, Colombia using the Collins and Paskewitz (1996) primers (n = 10, 546 bp). Only one haplotype was found. Primers sequences are underlined and in bold.

cause in Zavortink's key (1973), the presence or absence of scales in VII tergite in the abdomen of adults must be verified at one point, which is an external feature of the genitalia. Thus, the dichotomous key of Harrison et al. (2012) is useful for separating females of both species.

As mentioned above, for several years, there was evidence of the presence of *An. pholidotus* rather than *An. lepidotus* in the Cunday-Villarrica focal region (González & Carrejo 2009). Even just three years after the report of Quiñones et al. (1984), the presence of *An. lepidotus* in that zone was questioned (Harrison et al. 2012).

The limited availability of specimens collected from the area for morphological analysis, the scarcity of associated stages, the poor condition of some of the samples and the difficulty of accessing the area to collect new biological information most likely delayed the opportunity to solve this problem by three decades.

The collections of mosquitoes for this study, which occurred between February 2009–October 2011, provided updated information about the species of anophelines in these focal areas. According to the morphological features of females, the larvae chaetotaxy, the male genitalia presented in the specimens of Cunday-Villarrica malaria focal area in Tolima and the analysis of the *COI* and *ITS2* sequences, *An. pholidotus* was the most common

species found and was most likely misidentified as *An. lepidotus* in the previous report (Quiñones et al. 1984).

It is now possible to differentiate females of *An. pholidotus* from *An. lepidotus* using the dichotomous key. The fact that females of *An. lepidotus* were not found by analysing 1,737 individual mosquitoes captured indicates that the species that was previously reported as the vector of malaria in the area since 1984 (Quiñones et al. 1984) was *An. pholidotus* and not *An. lepidotus*.

The sequences obtained in this investigation, both *COI* and *ITS2*, showed that the tested specimens correspond to *An. pholidotus*, thus confirming the results of the morphology analysis. Sequences from three localities of Tolima exhibit no intraspecific variability, showing that *An. pholidotus* is present in the study area as a single species, although this may not be true in other parts of the country due to the high altitude of certain mountain ranges. Olano et al. (2001) and González and Carrejo (2009) provided records of supposed "*An. lepidotus*" in different departments of the country. For example, we have identified, by morphology, three females from Valle del Cauca (PNN Farallones: Cali, Pance, VI-6-1984, 1,500 m) as *An. pholidotus*. This result indicates that there is a possibility that *An. pholidotus* has a wider distribution than previously thought and that many of the specimens identified as *An. lepidotus* may correspond to *An. pholidotus*. Moreover, the sequences mentioned above show 100% homology with sequences of *An. pholidotus* from Táchira, Venezuela, recorded in GenBank by Harrison et al. (2012). As a result, it is important to determine whether this homology is generalised to all countries where *An. pholidotus* is present.

Harrison et al. (2012) inadvertently forgot to include the precise location of the record for *An. pholidotus* in Costa Rica. A single female in the mosquito collection located at the NMNH at the Smithsonian Institution was collected resting on the wall of a highway tunnel. The collection data record was as follows: Costa Rica: Cartago, Res. Tapanti; unnamed tribs., Ca. 9 Km (road NW tunnel, 9.72 N, 83.78 W, 8.9.vi.1988, elev. 1,400 m, CM & OS Flint, Molzenthal).

One of the key aspects for optimising control measures is an appropriate taxonomic determination of the species that are present and those that are responsible for transmission. Taxonomic studies in *Kerteszia* species are relevant because the distribution of this subgenus along the continent is still poorly known, mainly due to the lack of research studies. Furthermore, the association of these species with areas where bromeliads are normally present means that environmental changes and human activity may restrict the distribution for some species in the subgenus (Marrelli et al. 2007). However, for other species, the same factors may stimulate the proliferation of breeding sites and possibly extend their geographical distribution (Solarte et al. 1994).

## REFERENCES

- Benítez JA, Rodríguez A, Sojo M, Lobo H, Villegas C, Oviedo L, Brown E 2004. Descripción de un brote epidémico de malaria de altura en un área originalmente sin malaria del estado de Trujillo, Venezuela. *Boletín de Malariología y Salud Ambiental* 44: 93-99.

- Carvajal H, de Herrera MA, Quintero J, Alzate A, Herrera S 1989. *Anopheles neivai*: a vector of malaria in the Pacific low lands of Colombia. *Trans R Soc Trop Med Hyg* 83: 609.
- Collins FH, Paskewitz SM 1996. A review of the use of ribosomal DNA (rDNA) to differentiate among cryptic *Anopheles* species. *Insect Mol Biol* 5: 1-9.
- Collucci E, Sallum MAM 2003. Phylogenetic analysis of the subgenus *Kerteszia* of *Anopheles* (Diptera: Culicidae: Anophelinae) based on morphological characters. *Insect Syst Evol* 34: 361-372.
- Cova-García P 1961. *Notas sobre los anofelinos de Venezuela y su identificación*, 2nd ed., Grafos, Caracas, 213 pp.
- de Carvalho-Pinto CJ, Lourenço-de-Oliveira R 2004. Isoenzimatic analysis of four *Anopheles* (*Kerteszia*) *cruzii* (Diptera: Culicidae) populations of Brazil. *Mem Inst Oswaldo Cruz* 99: 471-475.
- Escobar J, González R, Quiñones ML, Wilkerson RC, Harrison B 2010. Presence of *Anopheles* (*Kerteszia*) *pholidotus* in a malaria focus in Colombia. In G Clark, Y Rubio-Palis, *Mosquito vector biology and control in Latin America - A 20th symposium*. *J Am Mosq Control Assoc* 26: 306-320.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3: 294-299.
- Forattini OP, Kakitani I, dos Santos RC, Ueno HM, Kobayashi KM 1999. Role of *Anopheles* (*Kerteszia*) *bellator* as malaria vector in southeastern Brazil (Diptera: Culicidae). *Mem Inst Oswaldo Cruz* 94: 715-718.
- González R, Carrejo N 2009. *Introducción al estudio taxonómico de Anopheles de Colombia. Claves y notas de distribución*, 2nd ed., Universidad del Valle, Cali, 248 pp.
- Gutierrez LA, Nelson N, Jaramillo LM, Muskus C, Luckhart S, Conn JE, Correa M 2008. Natural infectivity of *Anopheles* species from the Pacific and Atlantic Regions of Colombia. *Acta Trop* 107: 99-105.
- Harbach RE, Navarro JC 1996. A new species of *Anopheles*, subgenus *Kerteszia* (Diptera: Culicidae) from Venezuela. *Ent Scand* 27: 207-216.
- Harrison BA, Ruiz-López F, Calderón G, Savage HM, Pecor JE, Wilkerson RC 2012. *Anopheles* (*Kerteszia*) *lepidotus* (Diptera: Culicidae), not the malaria vector we thought it was: revised male and female morphology; larva, pupa and male genitalia characters and molecular verification. *Zootaxa* 3218: 1-17.
- Komp WHW 1937. The species of the subgenus *Kerteszia* of *Anopheles*. *Ann Entomol Soc Am* 30: 492-529.
- Linton Y-M, Harbach RE, Anthony TG, Chang MS, Asmad M 2001. Morphological and molecular identity of *Anopheles* (*Cellia*) *sundaicus* (Diptera: Culicidae), the nominotypical member of a malaria vector species complex in Southeast Asia. *Syst Entomol* 26: 357-366.
- Maddison DR, Maddison WR 2003. *MacClade v.4.06: analysis of phylogeny and character evolution*, Sinauer Associates, Sunderland, MA.
- Marrelli MT, Malafronte RS, Sallum MAM, Natal D 2007. *Kerteszia* subgenus of *Anopheles* associated with the Brazilian Atlantic rainforest: current knowledge and future challenges. *Malar J* 6: 127.
- Montoya J, González R, Palma G, Solarte Y, Olano V 1994. Malaria. In BL Travi, J Montoya-Lerma, Centro Internacional de Entrenamiento e investigaciones Médicas (eds.), *Manual de entomología médica para investigadores de América Latina*, CI-DEIM, Cali, p. 163-209.
- Montoya-Lerma J, Solarte YA, Giraldo-Calderón GI, Quiñones ML, Ruiz-López F, Wilkerson RC, González R 2011. Malaria vector species in Colombia - A review. *Mem Inst Oswaldo Cruz* 106 (Suppl. I): 223-238.
- Olano V, Brochero H, Sáenz R, Quiñones M, Molina J 2001. Mapas preliminares de la distribución de *Anopheles* vectores de malaria en Colombia. *Biomédica* 21: 402-403.
- Quiñones M, Suárez M, Rodríguez A, Fleming G, Galvis L 1984. Comportamiento de *Anopheles* (*Kerteszia*) *lepidotus* Zavortink, 1973, y su incriminación como posible vector de malaria en el Departamento del Tolima, Colombia. *Biomédica* 4: 5-13.
- Rubio-Palis Y, Zimmerman RH 1997. Ecoregional classification of malaria vectors in the Neotropics. *J Med Entomol* 34: 499-510.
- Ruiz F, Linton Y-M, Ponsonby DJ, Conn JE, Herrera M, Quiñones ML, Vélez ID, Wilkerson RC 2010. Molecular comparison of topotypic specimens confirms *Anopheles* (*Nyssorhynchus*) *dunhami* Causey (Diptera: Culicidae) in the Colombian Amazon. *Mem Inst Oswaldo Cruz* 105: 899-903.
- Sallum MAM, Schultz TR, Foster PG, Wirtz RA, Wilkerson RC 2002. Phylogeny of Anophelinae (Diptera: Culicidae) based on nuclear ribosomal and mitochondrial DNA sequences. *Syst Entomol* 27: 361-381.
- Solarte Y, González R, Hurtado JC, Alzate A 1994. Influencia de la vegetación en la presencia, distribución y abundancia de criaderos de tres especies de *Anopheles* (DIP.: Culicidae) en la parte baja del río Naya, costa pacífica de Colombia. *Bol Mus Ent Univ Valle* 2: 55-71.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S 2011. MEGA 5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance and maximum parsimony methods. *Mol Biol Evol* 28: 2731-2739.
- Ueno HM, Forattini OP, Kakitani I 2007. Vertical and seasonal distribution of *Anopheles* (*Kerteszia*) in Ilha Comprida, southeastern Brazil. *Rev Saude Publica* 41: 269-275.
- Zavortink TJ 1973. Mosquitoes studies (Diptera: Culicidae). XXIX - A review of the subgenus *Kerteszia* of *Anopheles*. *Contrib Amer Ent Inst* 9: 1-59.