

Frequency and molecular characterisation of *Entamoeba histolytica*, *Entamoeba dispar*, *Entamoeba moshkovskii*, and *Entamoeba hartmanni* in the context of water scarcity in northeastern Brazil

Deiviane Aparecida Calegar¹, Beatriz Coronato Nunes², Kerla Joeline Lima Monteiro²,
Jéssica Pereira dos Santos³, Helena Keiko Toma⁴, Tais Ferreira Gomes⁵,
Marli Maria Lima⁵, Márcio Neves Bóia¹, Filipe Anibal Carvalho-Costa^{2,3/+}

¹Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Biologia e Parasitologia de Mamíferos Silvestres Reservatórios, Rio de Janeiro, RJ, Brasil ²Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Epidemiologia e Sistemática Molecular, Rio de Janeiro, RJ, Brasil ³Fundação Oswaldo Cruz, Escritório Técnico Regional Fiocruz Piauí, Brasil

⁴Universidade Federal do Rio de Janeiro, Laboratório de Diagnóstico Molecular e Hematologia, Rio de Janeiro, RJ, Brasil

⁵Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Ecoepidemiologia da Doença de Chagas, Rio de Janeiro, RJ, Brasil

This study aimed to estimate the frequency, associated factors, and molecular characterisation of Entamoeba histolytica, Entamoeba dispar, Entamoeba moshkovskii, and Entamoeba hartmanni infections. We performed a survey (n = 213 subjects) to obtain parasitological, sanitation, and sociodemographic data. Faecal samples were processed through flotation and centrifugation methods. E. histolytica, E. dispar, E. moshkovskii, and E. hartmanni were identified by nested-polymerase chain reaction (PCR). The overall prevalence of infection was 22/213 (10.3%). The infection rate among subjects who drink rainwater collected from roofs in tanks was higher than the rate in subjects who drink desalinated water pumped from wells; similarly, the infection rate among subjects who practice open defecation was significantly higher than that of subjects with latrines. Out of the 22 samples positive for morphologically indistinguishable Entamoeba species, the differentiation by PCR was successful for 21. The species distribution was as follows: 57.1% to E. dispar, 23.8% to E. histolytica, 14.3% to E. histolytica and E. dispar, and 4.8% E. dispar and E. hartmanni. These data suggest a high prevalence of asymptomatic infection by the group of morphologically indistinguishable Entamoeba histolytica/dispar/moshkovskii complex and E. hartmanni species. In this context of water scarcity, the sanitary and socioenvironmental characteristics of the region appear to favour transmission.

Key words: parasites - protozoa - entamoebiasis - epidemiology - Brazil

Intestinal protozoan infections are closely related to a lack of proper sanitation and environmental contamination with faecal matter. Thus, their prevalence is higher in specific environmental scenarios that occur most often in developing countries (Ojha et al. 2014, Turkeltaub et al. 2015). Amoebiasis is a potentially severe and life threatening infection caused by enteric protozoa (Ralston & Petri Jr 2011, Skappak et al. 2014), most commonly *Entamoeba histolytica*, which is distributed worldwide (WHO 1997, Jackson 1998). The motile (trophozoite) form of *E. histolytica* inhabits the human colon where it multiplies and differentiates into cysts that are released into the environment. In turn, these cysts are responsible for transmitting the infection to another host via the faecal-oral route. The parasite invades the intestinal mucosa and causes many forms of invasive disease, including dysentery (Lin & Kao 2013). The parasite also exhibits bloodborne spreading and causes extraintestinal lesions, mainly liver

abscesses (Wuerz et al. 2012). The latter form occurs only rarely. Invasive disease occurs when virulent trophozoites disrupt the mucopithelial barrier by crossing the mucus layer, thereby damaging intestinal cells. This damage leads to inflammation and, consequently, dysentery (Thibeaux et al. 2013). Nevertheless, the majority of infections seem to be asymptomatic (Chacín-Bonilla 2013).

The existence of nonpathogenic indistinguishable *E. histolytica/Entamoeba dispar/Entamoeba moshkovskii* complex and *Entamoeba hartmanni* organisms capable of inhabiting the human intestine as commensals has been recognised for many decades. For instance, in 1926, Brumpt proposed the existence of *E. dispar*, a species indistinguishable by light microscopy from *E. histolytica*. However, *E. dispar* exhibits distinct physiological, biochemical, and ultrastructural characteristics, the latter of which have been described more recently (Goldman 1969, Jackson 1998, Pimenta et al. 2002). Another four-nucleated morphologically identical organism, *E. moshkovskii*, has been observed in sewage as a free-living amoeba, but is also capable of colonising the human intestine (Tshalaia 1941, Ngui et al. 2012). In addition, differential diagnosis should also consider the nonpathogenic species *E. hartmanni*, which can be distinguished from *E. histolytica* by its small cyst size (5-10 µm in diameter). In contrast, the diameter of *E. histolytica* cysts ranges from 12-14 µm (Brumpt 1949).

doi: 10.1590/0074-02760150383

DAC and BCN contributed equally to this study.

+ Corresponding author: guaritiba@ioc.fiocruz.br

Received 7 October 2015

Accepted 4 January 2016

More recently, dysentery and extraintestinal disease have been proposed to be potentially associated with *E. dispar* and *E. moshkovskii* (Parija & Khairnar 2005, Costa et al. 2010). These findings complicated our understanding of the pathogenic behaviour and public health importance of indistinguishable *E. histolytica*/*E. dispar*/*E. moshkovskii* complex and *E. hartmanni* parasites (Oliveira et al. 2015).

Vast rural areas in northeastern Brazil are characterised by deficits in sanitation infrastructure. Moreover, improper disposal of waste occurs frequently. These semiarid regions are also subjected to water stress due to prolonged droughts. Therefore, alternative water management approaches have been applied in this region (Rasella 2013). In this context, specific epidemiological scenarios associated with water scarcity could favour transmission of enteric pathogens. For example, water must be stored for many months during the dry season.

This study aimed to use molecular techniques to estimate the frequencies of infection with *E. histolytica*, *E. dispar*, *E. moshkovskii*, and *E. hartmanni* in a population subjected to water scarcity in the Northeast Region of Brazil. This study also aimed to identify factors associated with these infections.

SUBJECTS, MATERIALS AND METHODS

Study area and population - This study was performed in Russas, a municipality located 165 km from Fortaleza, the capital of the state of Ceará (Fig. 1). This region belongs to the semiarid region of northeastern Brazil, in the *Caatinga* biome. Russas has 74,243 inhabitants and a total area of 1,588 km². The study included four rural communities in the municipality: Riacho do Barro (132 inhabitants), Timbaúba do Pitingão (109 inhabitants), Barracão (315 inhabitants), and Patos de Tito (54 inhabitants). Russas has a hot, dry climate and is subjected to prolonged droughts. The rainy season typically extends from December-June (annual rainfall in 2013 = 418 mm, mean annual rainfall = 792.6 mm). Nev-

ertheless, seasonal rains have been reduced in the last few years and the region has been subjected to severe drought during the field work periods.

Study design and sampling strategy - We performed a cross-sectional survey from August-September 2013. The survey included 213 subjects (70 families): 53 subjects (18 families) from Timbaúba do Pitingão, 28 subjects (9 families) from Riacho do Barro, 119 subjects (38 families) from Barracão and 13 subjects (5 families) from Patos do Tito. Therefore, our study included 35% of the 610 residents in the four communities. We designed our sampling strategy specifically to include all households with children. During domicile visits, researchers distributed bottles without preservatives for faeces collection and obtained sanitation and sociodemographic data. In addition, the field team investigated whether the residents presented symptoms consistent with amoebiasis, such as diarrhoea, presence of mucus, pus, and/or blood in the stool, and abdominal pain, among others. The baseline characteristics of the study subjects are presented in Table I. Stool samples were collected the next day at each household and were transported to the field laboratory under refrigeration (4°C). The rates of *E. histolytica*, *E. dispar*, *E. moshkovskii*, and *E. hartmanni* detection in distinct sociodemographic settings were compared using Fisher's exact test. Statistical significance was established at $p < 0.05$.

Laboratory procedures - Initially, faecal samples were processed through the zinc sulphate flotation (Faust technique) and the formalin-ethyl-acetate centrifugation (modified Ritchie technique) methods (Faust et al. 1938, Young et al. 1979). For the Faust technique, 7 mL of gauze-filtered faecal suspension was spun by centrifugation and the resultant pellet was re-suspended in zinc sulphate solution (1,180 g/mL). The suspension was shaken and spun by centrifugation again, after which the resultant supernatant was examined by light




Fig. 1: map of the study area (Russas, state of Ceará, Brazil, 2013).

TABLE I

Sociodemographic characteristics of the studied population, Russas, state of Ceará, Brazil, 2013

Characteristics	n (%)
Gender	
Male	106 (49.8)
Female	107 (50.2)
Age group (years)	
0-4	18 (8.4)
5-9	27 (12.7)
10-14	37 (17.4)
15-19	14 (6.6)
> 19	117 (54.9)
Community	
Barracão	119 (55.9)
Patos do Tito	13 (6.1)
Riacho do Barro	28 (13.1)
Timbaúba do Pitingão	53 (24.9)
Income strata	
Extreme poverty (< US\$ ^a 17)	20 (9.4)
Poverty (US\$ 17-34)	27 (12.7)
Not poor (> US\$ 34)	166 (77.9)
Source of drinking water	
Desalinated brackish water from wells	138 (64.8)
Rain water stored in cisterns	56 (26.3)
Other	19 (8.9)
Sanitation facilities	
Latrine	166 (77.9)
Open defecation	47 (22.1)

^a: US\$ 1.00 = R\$ 4,00 (22 September 2015).

microscopy. For the Ritchie method, gauze-filtered faecal suspensions were spun by centrifugation and the resultant pellets were re-suspended in 5 mL of water and 3 mL of ethyl-acetate was added to each suspension. The sedimented matter was examined by light microscopy. It was not possible to perform permanent smear staining for light microscopy or to measure amoebae cysts in the field laboratory; thus, *E. hartmanni* could not be distinguished from *E. histolytica*, *E. dispar*, and *E. moshkovskii*. Faecal samples were cryopreserved and transported to the city of Rio de Janeiro, Brazil for molecular tests. All indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni* positive faecal samples were subjected to DNA extraction using the ZR Fungal/Bacterial DNA MiniPrep™ kit. Nested-polymerase chain reaction (PCR) was performed according to the protocol described by Paglia and Visca (2004). Initially, 1,076 bp fragment of the small subunit rRNA gene sequence common to the *Entamoeba* genus was amplified using primers E1 (5-TGCTGTGATTA-AAACGCT-3) and E2 (5-TTAACATTTCAATCTC-GG-3). Nested-PCR was performed with primers Eh-L

(5-ACATTTTGAAGACTTTATGTAAGTA-3) and Eh-R (5-CAGATCTAGAAACAATGCTTCTCT-3), which are specific for *E. histolytica* and amplify a 427 bp fragment, Ed-L (5-GTTAGTTATCTAATTTTCGATTAGAA-3) and Ed-R (5-ACACCACTTACTATCCCTACC-3), which are specific for *E. dispar* and amplify a 195 bp product, and Mos 1 (5-GAAACCAAGAGTTTCACAAC-3) and Mos 2 (5-CAATATAAGGCTTGGATGAT-3), which are specific for *E. moshkovskii* and yield a 553 bp product (Paglia & Visca 2004, Lau et al. 2013). Molecular characterisation of *E. hartmanni* was performed essentially as described by Gomes et al. (2014), but with minor modifications. Briefly, primers EhartR1 mod (5-ATTGTCTTCACTATTCCATGCC-3) and EhartF mod (5-CCAGCTTTCCAAACATGATG-3) were used to amplify a 186 bp product. PCR products were resolved on 1.5% agarose gels, stained with ethidium bromide, and visualised *via* ultraviolet illumination.

Ethics - This study was approved by the Ethical Committee in Research with Humans, Oswaldo Cruz Institute, Oswaldo Cruz Foundation (CAAE: 12125713.5.0000.5248).

RESULTS

The overall prevalence of infection with indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni* organisms was 22/213 (10.3%). Of these 22 positive faecal samples, one was identified only through the flotation (Faust) method, 13 were identified only with the centrifugation (Ritchie 1948) method, and eight were identified with both techniques. The detection rates of nonpathogenic amoebas were as follows: *Endolimax nana*, 4.2% (n = 9), *Entamoeba coli*, 11.3% (n = 24), and *Iodamoeba butschlii*, 7% (n = 15). *Giardia intestinalis* was detected in 30 subjects (14.1%). The age distribution of indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni* infections is presented in Fig. 2. Regarding infection positivity according to sex, indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni* infections were found in 12/106 males and 10/107 females (p = 0.704).

As presented in Table II, the detection rate of indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni* among subjects who drink rain-water collected from roofs in tanks was higher than the rate in people who drink desalinated water pumped from wells. In addition, the detection rate among subjects who practice open defecation was significantly higher than that of inhabitants who have latrines. The positivity rates of subjects in different income strata were similar.

Species-level identification could be performed for 21 of the 22 samples positive for indistinguishable *E. histolytica/E. dispar/E. moshkovskii* complex and *E. hartmanni*. The species distribution was as follows: 12 (57.1%) *E. dispar*, 5 (23.8%) *E. histolytica*, 3 (14.3%) co-infections with *E. histolytica* and *E. dispar*, and one (4.8%) co-infection with *E. dispar* and *E. hartmanni* (Fig. 3). No sample was positive for *E. moshkovskii*. The age distributions of subjects infected with different species are shown in Fig. 4.




Fig. 2: results of parasitological analysis for *Entamoeba histolytica*/*Entamoeba dispar*/*Entamoeba moshkovskii* complex and *Entamoeba hartmanni* considering the age groups (Russas, state of Ceará, Brazil, 2013).

TABLE II

Rate of detection of *Entamoeba histolytica*/*Entamoeba dispar*/*Entamoeba moshkovskii* complex and *Entamoeba hartmanni* by source of drinking water, place of defecation, and income, Russas, state of Ceará, Brazil, 2013

	Positive/tested subjects (% of positive)	p ^a
Source of drinking water		
Desalinated brackish water from wells	9/138 (6.5)	0.054
Rain water stored in cisterns	9/56 (16.1)	
Sanitation facilities		
Latrine	13/166 (7.8)	0.032
Open defecation	9/47 (19.1)	
Family month income per capita		
< US\$ ^b 17	4/20 (20)	0.308
US\$ 17-34	2/27 (7.4)	
> US\$ 34	16/166 (9.6)	

a: Fisher exact test; b: US\$ 1.00 = R\$ 4,00 (22 September 2015).

DISCUSSION

A key issue for understanding the morbidity associated with amoebiasis is to define the proportion of infections associated with the pathogenic species *E. histolytica*. Interestingly, studies in different regions have shown that many subjects infected with indistinguishable *E. histolytica*/*E. dispar*/*E. moshkovskii* complex and *E. hartmanni* parasites actually harbour low-pathogenicity species such as *E. dispar*, *E. moshkovskii*, or even *E. hartmanni* (Gomes et al. 2014, Nair & Variyam 2014, Efunshile et al. 2015, Nath et al. 2015). The proportions of these subjects are variable, but can be quite high.

E. dispar and *E. moshkovskii* are indistinguishable from *E. histolytica* by light microscopy. Thus, routine parasitological techniques are not suitable for discriminating these organisms. This limitation means that a

significant number of patients being treated with anti-parasitic drugs such as metronidazole may not actually be infected with *E. histolytica*.

In the present study, approximately two-thirds of all infections were not caused by *E. histolytica*. We note that all subjects were asymptomatic at the time of the stool test. Even so, we infer that nonpathogenic species are detected more frequently than *E. histolytica* in the studied area. This observation is particularly relevant because increasing importance has been given to traditionally nonpathogenic species such as *E. dispar* and *E. moshkovskii*, since invasive amoebiasis has been demonstrated to be associated with these species (Parija & Khairnar 2005). It is likely that the determinants of invasive amoebiasis are complex and also involve host factors (Bosch & Siderovski 2013, Thibaux et al. 2013).




Fig. 3: detection and differentiation of *Entamoeba histolytica*, *Entamoeba moshkovskii*, *Entamoeba dispar* and *Entamoeba hartmanni* by nested-polymerase chain reaction. PCR products were visualised in 1.5% agarose gel with EtBr staining. Line 1: 100 bp DNA ladder; 2, 3: one faecal sample with mixed infection by *E. dispar* and *E. histolytica*, respectively; 4, 6, 7: faecal samples positive for *E. dispar*; 5, 14: empty wells; 8, 9: faecal samples positive for *E. histolytica*; 10: negative control for *E. dispar*; 11: negative control for *E. histolytica*; 12: positive control for *E. dispar*; 13: positive control for *E. histolytica*; 15: faecal sample positive for *E. hartmanni*; 16: negative control.




Fig. 4: frequency of identification of *Entamoeba histolytica*, *Entamoeba moshkovskii*, *Entamoeba dispar* and *Entamoeba hartmanni* by nested-polymerase chain reaction by age group in 21 positive subjects (Russas, state of Ceará, Brazil, 2013).

The nonpathogenic species *E. hartmanni* can be distinguished from *E. histolytica*, *E. dispar*, and *E. moshkovskii* by light microscopy. However, this distinction requires detailed observation of nuclear structures, which requires permanent smear staining, an ocular micrometer, and a highly skilled parasitologist. These criteria are hard to meet for many laboratories. We propose that the possibility of *E. hartmanni* infection should also be considered in people who excrete indistinguishable *E. histolytica*/*E. dispar*/*E. moshkovskii* complex and *E. hartmanni* cysts. In the present study, *E. hartmanni* was detected in one of the indistinguishable *E. histolytica*/*E. dispar*/*E. moshkovskii* complex and *E. hartmanni* positive samples.

The study population is located in a sociodemographic and environmental setting characterised by deficits in sanitation infrastructure and water stress. The study area is located in a low-rainfall region in the *Caatinga* biome that is subjected to prolonged droughts and prone to desertification. Nonpotable water is obtained from a reservoir in the locality and used for livestock watering and other suitable applications. In the last decade, a strategy has been implemented in which rainwater is collected during the rainy season from roofs *via* gutters. This collected rainwater is stored in household tanks for later use during droughts. This strategy has significantly improved access to drinking water in the study area. Artesian wells constructed in the region are another source of drinking water. However, this water is brackish and must be desalinated before consumption. We found that the rate of *E. histolytica*, *E. dispar*, and *E. hartmanni* positivity was almost three times higher in subjects who drink collected rainwater than in subjects who drink desalinated brackish water drawn from the artesian wells. We hypothesise that the long period (between the dry season and the rainy season) of rainwater storage in tanks favours contamination with amoeba cysts, thereby enabling transmission. Interestingly, consumption of rainwater captured from roofs has been demonstrated to reduce the prevalence of *G. intestinalis* infection in a semiarid region in northeastern Brazil (Fonseca et al. 2014). Regarding the place of defecation, subjects who practice open defecation exhibited a significantly higher positive rate compared with subjects who defecate in latrines. Moreover, an even higher positive rate was observed in people who deposit faeces directly into the soil compared with subjects with rudimentary tanks.

In some regions of the world, including Latin America, inadequate sanitary conditions facilitate the transmission of amoebiasis, thereby generating high prevalence rates (Braga et al. 1998, Ramos et al. 2005). In these scenarios, invasive amebic dysentery and liver abscesses are expected to occur. However, these diseases were not observed in the present study. Severe cases of amoebiasis are identified infrequently in Brazil, which may be explained by the relative improvement of living conditions over the past few decades.

Cumulatively, our data suggest a high prevalence of asymptomatic infection with indistinguishable *E. histolytica*/*E. dispar*/*E. moshkovskii* complex and *E. hartmanni* parasites. These asymptomatic infections appear to be caused by predominantly nonpathogenic species or parasites with low pathogenic potential. In the context of scarce water resources, the sanitary and socioenvironmental characteristics of the region appear to be associated with transmission.

REFERENCES

- Bosch DE, Siderovski DP 2013. G protein signaling in the parasite *Entamoeba histolytica*. *Exp Mol Med* 45: e15.
- Braga LL, Mendonça Y, Paiva CA, Sales A, Cavalcante AL, Mann BJ 1998. Seropositivity for and intestinal colonization with *Entamoeba histolytica* and *Entamoeba dispar* in individuals in northeastern Brazil. *J Clin Microbiol* 10: 3044-3045.

- Brumpt E 1926. Individualité de l'Entamoeba dispar. Présentation de pièces. *Bull Soc Pathol Exot* 19: 399-404.
- Brumpt E 1949. *Précis de parasitologie*, Masson & Cie, Paris, 2138 pp.
- Chacín-Bonilla L 2013. Amebiasis: aspectos clínicos, terapéuticos y de diagnóstico de la infección. *Rev Med Chil* 5: 609-615.
- Costa CA, Brito KN, Gomes MA, Caliarí MV 2010. Histopathological and immunohistochemical study of the hepatic lesions experimentally induced by *Entamoeba dispar*. *Eur J Histochem* 3: e39.
- Efunshile MA, Ngwu BA, Kurtzhals JA, Sahar S, König B, Stensvold CR 2015. Molecular detection of the carriage rate of four intestinal protozoa with real-time polymerase chain reaction: possible overdiagnosis of *Entamoeba histolytica* in Nigeria. *Am J Trop Med Hyg* 93: 257-262.
- Faust EC, D'antoni JS, Odom V, Miller MJ, Peres C, Sawitz W, Thomen LF, Tobie J, Walker JH 1938. A critical study of clinical laboratory techniques of the diagnosis of protozoan cysts and helminth eggs in feces. I - Preliminary communication. *Am J Trop Med Hyg* 18: 169-183.
- Fonseca JE, Carneiro M, Pena JL, Colosimo EA, da Silva NB, da Costa AG, Moreira LE, Cairncross S, Heller L 2014. Reducing occurrence of *Giardia duodenalis* in children living in semiarid regions: impact of a large scale rainwater harvesting initiative. *PLoS Negl Trop Dis* 6: 2943.
- Goldman M 1969. *Entamoeba histolytica*-like amoebae occurring in man. *Bull World Health Organ* 3: 355.
- Gomes TDS, Garcia MC, Cunha FDS, de Macedo HW, Peralta JM, Peralta RHS 2014. Differential diagnosis of *Entamoeba* spp in clinical stool samples using SYBR green real-time polymerase chain reaction. *Sci World J* 2014: 8 pp.
- Jackson TE 1998. *Entamoeba histolytica* and *Entamoeba dispar* are distinct species; clinical, epidemiological, and serological evidence. *Int J Parasitol* 28: 181-186.
- Lau YL, Anthony C, Fakhrrrazi SA, Ibrahim J, Ithoi I, Mahmud R 2013. Real-time PCR assay in differentiating *Entamoeba histolytica*, *Entamoeba dispar*, and *Entamoeba moshkovskii* infections in Orang Asli settlements in Malaysia. *Parasit Vectors* 6: 250.
- Lin CC, Kao KY 2013. Ameboma: a colon carcinoma-like lesion in a colonoscopy finding. *Case Rep Gastroenterol* 7: 438-441.
- Nair GV, Variyam EP 2014. Noninvasive intestinal amebiasis: *Entamoeba histolytica* colonization without invasion. *Curr Opin Infect Dis* 5: 465-469.
- Nath J, Hussain G, Singha B, Paul J, Ghosh SK 2015. Burden of major diarrheagenic protozoan parasitic co-infection among amoebic dysentery cases from north east India: a case report. *Parasitol* 10: 1318-1325.
- Ngui R, Angal L, Fakhrrrazi SA, Lian YL, Ling LY, Ibrahim J, Mahmud R 2012 Differentiating *Entamoeba histolytica*, *Entamoeba dispar*, and *Entamoeba moshkovskii* using nested polymerase chain reaction (PCR) in rural communities in Malaysia. *Parasit Vectors* 5: 187.
- Ojha SC, Jaide C, Jinawath N, Rotjanapan P, Baral P 2014. Geohelminths: public health significance. *J Infect Dev Ctries* 8: 5-16.
- Oliveira FMS, Neumann E, Gomes MA, Caliarí MV 2015. *Entamoeba dispar*: could it be pathogenic. *Trop Parasitol* 10: e0137327.
- Paglia MG, Visca P 2004. An improved PCR-based method for detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* in formalin-fixed stools. *Acta Trop* 92: 273-277.
- Parija SC, Khairnar K 2005. *Entamoeba moshkovskii* and *Entamoeba dispar*-associated infections in Pondicherry, India. *J Health Popul Nutr* 23: 292-295.
- Pimenta PF, Diamond LS, Mirelman D 2002. *Entamoeba histolytica* Schaudinn, 1903 and *Entamoeba dispar* Brumpt, 1925: differences in their cell surfaces and in the bacteria-containing vacuoles. *J Euk Microbiol* 49: 209-219.
- Ralston KS, Petri Jr WA 2011. Tissue destruction and invasion by *Entamoeba histolytica*. *Trends Parasitol* 27: 253-263.
- Ramos F, Moran P, González E, García G, Ramiro M, Gómez A, León MDCG, Melendro EI, Valadez A, Ximénez C 2005. High prevalence rate of *Entamoeba histolytica* asymptomatic infection in a rural Mexican community. *Am J Trop Med Hyg* 73: 87-91.
- Rasella D 2013. Impact of the water for all program (PAT) on childhood morbidity and mortality from diarrhea in the Bahia state, Brazil. *Cad Saude Publica* 29: 40-50.
- Ritchie L 1948. An ether sedimentation technique for routine stool examinations. *Bull US Army Med Dep* 8: 326.
- Skappak C, Akierman S, Belga S, Novak K, Chadee K, Urbanski SJ, Church D, Beck PL 2014. Invasive amoebiasis: a review of *Entamoeba* infections highlighted with case reports. *Can J Gastroenterol Hepatol* 28: 355-359.
- Thibeaux R, Weber C, Hon CC, Dillies MA, Avé P, Coppée JY, Labruyère E, Guillén N 2013. Identification of the virulence landscape essential for *Entamoeba histolytica* invasion of the human colon. *PLoS Pathog* 9: e1003824.
- Tshalaia LE 1941. On a species of *Entamoeba* detected in sewage effluents. *Med Parazitol (Mosk)* 10: 244-252.
- Turkeltaub JA, McCarty TR, Hotez PJ 2015. The intestinal protozoa: emerging impact on global health and development. *Curr Opin Gastroenterol* 31: 38-44.
- WHO - World Health Organization 1997. WHO/PAHO/UNESCO report. A consultation with experts on amoebiasis. *Epidemiol Bull* 18: 13-14.
- Wuerz T, Kane JB, Boggild AK, Krajden S, Keystone JS, Fuksa M, Kain KC, Warren R, Kempston J, Anderson J 2012. A review of amoebic liver abscess for clinicians in a nonendemic setting. *Can J Gastroenterol* 26: 729-733.
- Young KH, Bullock SL, Melvin DM, Spruill CL 1979. Ethyl acetate as a substitute for diethyl ether in the formalin-ether sedimentation technique. *J Clin Microbiol* 10: 852-853.