

LETTER TO THE EDITOR

Emergence of *Acinetobacter baumannii* ST730 carrying the *bla*_{OXA-72} gene in Brazil

Mariana Pagano^{1,2/+}, Franciéli P Rozales^{1,2},
Diego Bertolini², Lisiane Rocha³, Jorge LM
Sampaio³, Afonso L Barth^{1,2}, Andreza F Martins^{2,4}

¹Universidade Federal do Rio Grande do Sul, Faculdade de Farmácia, Programa de Pós-Graduação em Ciências Farmacêuticas, Porto Alegre, RS, Brasil ²Hospital de Clínicas de Porto Alegre, Centro de Pesquisa Experimental, Laboratório de Pesquisa em Resistência Bacteriana, Porto Alegre, RS, Brasil ³Laboratório Fleury, Porto Alegre, RS, Brasil ⁴Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Microbiologia Agrícola e do Ambiente, Porto Alegre, RS, Brasil

Over the last decade, *Acinetobacter baumannii* resistant to carbapenems has emerged in many medical centres and has been commonly associated with high morbimortality. In Brazil, this resistance is mainly attributed to the spread of OXA-23-producing clones and, to a lesser extent, to OXA-143-producing clones. Here, we describe, for the first time, two OXA-72-producing *A. baumannii* isolates in southern Brazil to a broad spectrum of antibiotics, except polymyxin B and tigecycline. Molecular typing by multilocus sequence typing (MLST) demonstrated that both OXA-72-producing isolates belong to a new sequence type (ST), ST730, which was recently identified in OXA-23-producing *A. baumannii* isolates in São Paulo, Brazil. We demonstrate that the two *A. baumannii* ST730 isolates carrying *bla*_{OXA-72} share a common ancestral origin with the *bla*_{OXA-23} producers in Brazil. This observation reinforces the importance of strain-typing methods in order to clarify the dynamics of the emergence of new clones in a geographic region.

Key words: *Acinetobacter baumannii* - MLST - OXA-72

Recently, a new sequence type, ST730, from an OXA-23-producing *Acinetobacter baumannii* has been deposited in the multilocus sequence typing (MLST) database (<http://www.pasteur.fr>) (Vasconcelos et al. 2015). In the present study, we describe two isolates of *A. baumannii* ST730 carrying the *bla*_{OXA-72} gene from different patients in a hospital located in southern Brazil.

In March 2013, a carbapenem-resistant *Acinetobacter* sp. was isolated using VITEK®2 system (bioMérieux, La Balme-les-Grottes, France) from the tracheal aspi-

rate (> 10⁶ CFU/mL) of a 76-year-old female ICU patient in a 312-bed tertiary care hospital in Porto Alegre, southern Brazil. A few days later, another carbapenem-resistance in *Acinetobacter* sp. was reported from a blood culture (positive in 12 h) of an 86-year-old male ICU patient admitted with sepsis in the same hospital. Both the isolates were identified as *A. baumannii* by matrix-assisted laser desorption/ionisation time-of-flight (MALDI-TOF; Bruker Daltonik, Bremen, Germany) spectrometry and *gyrB* multiplex polymerase chain reaction (PCR). Disc-diffusion assays demonstrated that the isolates were resistant to meropenem, imipenem, amikacin, ampicillin-sulbactam, cefepime and ceftazidime. However, the isolates were found to be susceptible to polymyxin (MICs 1 and 2 µg/mL) and tigecycline (MICs 0.5 and 0.75 µg/mL) as per microdilution and Etest® assays, respectively. The results obtained were interpreted according to the CLSI guidelines (CLSI 2014). The carbapenemase genes (*bla*_{OXA-23-like}, *bla*_{OXA-24/40-like}, *bla*_{OXA-58-like} and *bla*_{OXA-143-like}) were investigated by multiplex PCR as previously described (Higgins et al. 2010). The *bla*_{OXA-24/40-like} gene was detected in both the isolates. Sequencing of *bla*_{OXA-24} gene (ABI 3500 Genetic Analyzer; Applied Biosystems, Foster City, CA, United States) identified the variant *bla*_{OXA-72}, which displayed 100% identity to the original *bla*_{OXA-72} gene (GenBank accession number AY739646.1). Clonal diversity, investigated by repetitive-sequence-based PCR (REP-PCR), revealed an identical profile of the isolates (Bou et al. 2000). MLST was performed according to the Institut Pasteur scheme (<http://www.pasteur.fr>) and both the isolates were identified as ST730.

Resistance to carbapenems among *A. baumannii* isolates from Brazil has been mostly related to the production of OXA-23, followed by OXA-143 (Vasconcelos et al. 2015). In fact, OXA-72-producing *A. baumannii* isolates are still uncommon in Brazil. The first two cases of *A. baumannii* carrying *bla*_{OXA-72} gene were reported from São Paulo (Southeast Brazil) in 2011 (Antonio et al. 2011, Werneck et al. 2011). Two years later, this gene was reported in two other *A. baumannii* isolates from Recife (Northeast Brazil) (Cavalcanti et al. 2013). Recently, a surveillance study evaluated nine hospitals from five different states, representative of all the Brazilian regions, and described an inter-hospital dissemination of 10 *A. baumannii* isolates containing *bla*_{OXA-72} in São Paulo (Vasconcelos et al. 2015). These data highlight the possibility of the spread of this gene in the country. Furthermore (Werneck et al. 2011), reported the presence of *bla*_{OXA-72} gene inserted on a plasmid of ~ 86 kb, highlighting its potential for spread.

Here, we describe, for the first time, two *A. baumannii* isolates harbouring the *bla*_{OXA-72} gene isolated from Rio Grande do Sul, southern Brazil. These data point towards the increasing diversity of oxacillinases among the clinical isolates of *Acinetobacter* spp. in Brazil. Recently, Cayô et al. (2015) deposited an OXA-23-producing isolate, *A. baumannii* ST730, in the Institut Pasteur database. ST730 is a single-locus variation of ST79 (CC79), responsible for the spread of *bla*_{OXA-23} in Latin America (Chagas et al. 2014, Vasconcelos et al. 2015). It is worrisome that

doi: 10.1590/0074-02760160188

+ Corresponding author: mari.pag@hotmail.com

Received 2 May 2016

Accepted 27 June 2016

the emergent OXA-72-producing clones share the same phylogenetic origin with the clones harbouring *bla*_{OXA-23} gene, since the latter demonstrated a remarkable capacity of dissemination and maintenance along the years in Brazilian hospitals (Pagano et al. 2015). The results presented in this study highlight the importance of monitoring the spread of successful clones associated with the dissemination of *A. baumannii* carrying *bla*_{OXA} in Brazil by molecular epidemiology methods such as MLST.

REFERENCES

- Antonio CS, Neves PR, Medeiros M, Mamizuka EM, de Araujo MRE, Lincopan N. High prevalence of carbapenem-resistant *Acinetobacter baumannii* carrying the *bla*_{OXA-143} gene in Brazilian hospitals. *Antimicrob Agents Chemother*. 2011; 55(3): 1322-3.
- Bou G, Cervero G, Dominguez MA, Quereda C, Martinez-Beltran J. PCR-based DNA fingerprinting (REP-PCR, AP-PCR) and pulsed-field gel electrophoresis characterization of a nosocomial outbreak caused by imipenem- and meropenem-resistant *Acinetobacter baumannii*. *Clin Microbiol Infect*. 2000; 6(12): 635-3.
- Cavalcanti FLS, Almeida AC, Vilela MA, de Morais Junior MA, de Morais MM, Leal-Balbino TC. Emergence of extensively drug-resistant OXA-72-producing *Acinetobacter baumannii* in Recife, Brazil: risk of clonal dissemination? *Diagn Microbiol Infect Dis*. 2013; 77(30): 250-1.
- Chagas TP, Carvalho KR, Santos ICO, Carvalho-Assef AP, Asensi MD. Characterization of carbapenem-resistant *Acinetobacter baumannii* in Brazil (2008-2011): countrywide spread of OXA-23-producing clones (CC15 and CC79). *Diagn Microbiol Infect Dis*. 2014; 79(4): 468-72.
- CLSI - Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing, Twenty-Fourth Informational Supplement, CLSI document M100-S24. Wayne: CLSI; 2014.
- Higgins PG, Lehmann M, Seifert H. Inclusion of OXA-143 primers in a multiplex polymerase chain reaction (PCR) for genes encoding prevalent OXA carbapenemases in *Acinetobacter* spp. *Int J Antimicrob Agents*. 2010; 35(3): 305.
- Pagano M, Barin J, Martins AF, Zavascki AP. High endemic rates of OXA-23-producing carbapenem-resistant *Acinetobacter baumannii* isolates caused by the persistence of major clones in hospitals in a Brazilian city 5 years after an outbreak. *Infect Control Hosp Epidemiol*. 2015; 36(7): 860-2.
- Vasconcelos AT, Barth AL, Zavascki AP, Gales AC, Levin AS, Lucaravski BR, et al. The changing epidemiology of *Acinetobacter* spp. producing OXA carbapenemases causing bloodstream infections in Brazil: a BrasNet report. *Diagn Microbiol Infect Dis*. 2015; 83(4): 382-5.
- Werneck JS, Picao RC, Carvalhaes CG, Cardoso JP, Gales AC. OXA-72-producing *Acinetobacter baumannii* in Brazil: a case report. *J Antimicrob Chemother*. 2011; 66(2): 452-4.