

The relations between dietary antioxidant vitamins intake and oxidative stress in follicular fluid and ART outcomes

Ashraf Kazemi¹ Ph.D., Fatemeh Ramezanzadeh² M.D., Mohammad Hosein Nasr-Esfahani³ Ph.D.

1. Reproductive Health Department, Nursing and Midwifery Care Research Center, School of Nursing and Midwifery, Isfahan University of Medical Sciences, Isfahan, Iran.
2. Vali-e-Asr Reproductive Health Research Center, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran.
3. Reproductive Biomedicine Center, Royan Institute for Animal Biotechnology, ACECR, Department of Reproduction and Development, Isfahan, Iran.

Corresponding Author:

Ashraf Kazemi, Reproductive Health Department, Isfahan University of Medical Sciences, Hezarjerib Ave., Isfahan, Iran.
Email: kazemi@nm.mui.ac.ir
Tel: (+98) 31 37922933

Received: 11 December 2014

Revised: 1 June 2015

Accepted: 6 June 2015

Abstract

Background: Oxidative stress (OS) in the follicular environment may affect on oocyte competence and antioxidant vitamins may modify its effects.

Objective: This study was conducted to examine the effect of dietary intake of vitamin A, C and E on OS in follicular environment and assisted reproduction technology (ART) outcomes.

Materials and Methods: In this observational prospective study, the intake levels of vitamin A, C, and E were measured by validated food frequency questionnaire and Malondialdehyde and the total antioxidant capacity (TAC) levels of follicular fluid (FF) in 219 women undergoing ART were assessed. The number of retrieved oocytes, percentages of metaphase II (MII) stage oocytes, fertilization rate, and embryo quality were also determined.

Results: No significant association was found between vitamins intake levels and OS biomarkers, but the mean of TAC level in FF among women who received vitamin C greater than 75 mg/d was higher than women with lower intakes ($p < 0.05$). The ART parameters were not related to the vitamin E intake level, but the normal cleaved embryo rate was positively related to vitamin A ($p < 0.05$) and vitamin C ($p = 0.02$) intake levels. Also, the percentage of MII oocytes ($p = 0.02$) and the fertilization rate ($p < 0.05$) were related to the vitamin C intake level. The relation between the TAC level in FF and ART outcomes were not significant.

Conclusion: Current results indicated that high dietary intake of vitamin C would be followed by increasing the TAC level in FF and improving the oocyte competence, but this effect of vitamin C is not dependent of increasing of antioxidant defense in follicular environment.

Key words: Oxidative stress, Vitamin A, Vitamin C, Vitamin E, Assisted reproduction, Fertilization, Embryo quality.

Introduction

In vivo, oocyte maturation is influenced by the interplay of the biochemical state of the oocyte and its surrounding microenvironment (1). In an optimal condition the follicle constitutes a complex and regulates microenvironment in which the oocyte acquires developmental competence at ovulation (2, 3). This environment contains large numbers of macrophages, neutrophils, and metabolically active granulosa cells, which produce reactive oxygen species (4). These oxygen-derived molecules are intermediary products of physiologic cellular metabolism (5). Under normal conditions, these molecules are neutralized by a defense system consisting of enzymatic and non-enzymatic antioxidants (6). An overproduction

of the reactive oxygen species and/or a deficit or depletion of the scavenging capacity of antioxidant defense systems leads to oxidative stress (OS) and oxidative damage of key intracellular components (7).

Some studies have suggested that OS in antral follicle may induce follicular atresia (8, 9) and influences oocyte competence (10-13). Therefore, the efforts aimed at reducing OS in a follicular environment can be considered. Some evidence suggests that dietary intervention could be an effective strategy to reduce systemic OS and its cellular damages (14, 15). Antioxidants are important in maintaining the oxidant-antioxidant balance in blood and tissues. The antioxidants vitamins, such as vitamins A, C, and E are important in restoring or maintaining the redox balance in blood and tissues (16).

In addition, for the embryo to acquire developmental potential, it may be important that antioxidants are stored in the oocyte during its growth and maturation phases. Tola *et al.* suggested that high intracellular antioxidant activity in granulosa cells had a positive effect on oocyte competence for fertilization (17). As a result, dietary antioxidant availability due to a rich antioxidant dietary intake could have an impact on the level of the OS in follicular environment.

Much of the interest in improving the quality of in vivo and in-vitro-matured human oocytes has focused on the effects of antioxidant supplementation (18). While, the dietary sources of vitamin intake, in addition to supplying vitamin, could have other metabolic effects (19). The effects of dietary antioxidant-rich intake on OS in the follicular environment have scarcely been investigated; besides diets rich with antioxidants deserve consideration in the context of the OS in follicular environment. The previous findings on the effect of ascorbic acid upon reproduction were contradictory. While, some studies in animal models showed the positive effects of ascorbic acid on the oocyte competence (20, 21), the results of other studies have not confirmed these findings (22). Also, some documents have shown the positive effect of using supplemental vitamin A on ART outcomes (23, 24). But there is no evidence to show that the positive effects of this vitamin on reproduction looking for antioxidant characteristics of vitamin A.

Therefore, the relationship between the dietary antioxidant vitamins intake level and lipid peroxidation product in FF, in addition, the relationship between the dietary antioxidant vitamin intake and assisted reproductive parameters were evaluated.

Materials and methods

For this observational prospective study, 219 non-donor in vitro fertilization/intracytoplasmic sperm insemination (IVF/ICSI) cycles were randomly conducted from July 2010 to April 2011 at Isfahan Fertility and Infertility centre. The study was approved by the Institutional Review Board and the Ethics Committee of Isfahan University of Medical Sciences. The exclusion criteria were male factor infertility, according to world health

organization (WHO) criteria (25), known systemic disease and considerable change in dietary pattern during assisted reproduction cycle. All eligible women gave written informed consent.

Assessments

At the start of ART cycle a validated semi-quantitative food frequency questionnaire (FFQ) including 168 food items (26) was used to measure calorie intake and vitamin intake, including vitamin A, C and E over the previous three months.

For all the main food items in the FFQ, the frequency per day was multiplied by the amount consumed, depending on the portion size, to compute the total amount consumed per day. The United States department of agriculture food composition table was used for most items. Depending on the appropriate vitamin intake, the subjects were divided into two groups for each vitamin.

For lifestyle variables, the participants were asked about smoking status (passive and active smoking) and about vitamin supplement (use Yes/No).

Assisted reproduction

Ovarian hyperstimulation using long protocol, involving gonadotropin releasing hormone agonist (Suprefact, Hoechst, Germany) and recombinant follicle stimulating hormone (Gonal F, Serono, Rockland, MA, USA, Follistim, Organon, Roseland, NJ) and/or human menopausal gonadotropin (Menogon, Ferring, Suffern, NY) administration were done and follicular maturation was monitored by transvaginal ultrasound examination.

Ovulation was triggered with 10,000 IU human chorionic gonadotropin (Pregnyl, Organon), when dominant follicles reach a follicular size of around 18mm. 36 hours later, the oocytes were retrieved transvaginally and used for IVF or ICSI procedure. After retrieving oocytes, 2 ml follicular fluid from one to 5 follicles (with ≤ 16 mm diameter) was pooled and centrifuged at 300 rpm for 17 min. of the supernatant was stored at -70°C . Maximum storage time was two weeks. Bloody samples and samples without oocyte were discarded.

Despite exclusion criteria of male factors infertility, as a routine protocol of Isfahan

Fertility and Infertility center, some of oocytes were inseminated through ICSI procedure.

Oocytes were considered fertilized when two pronuclei was observed 17-19h post sperm insemination. According to Veeck's criteria high-quality embryos at 3 days after oocyte retrieval were defined as grades one and two (27). The percentage of embryo with good quality was calculated by dividing the number of high-quality embryos by the number of total embryos. Embryos with more than 5 cells on 3 days post IVF/ICSI were considered as normal cleaved embryos and the normal cleaved embryos rate was calculated by dividing the number of the normal cleaved embryos by the number of total embryos. In ICSI cycles, the percentages of the MII stage were calculated by dividing the number of oocytes with first polar body divided by the number of assessed oocytes. The data about the number of used gonadotropin were recorded based on the patients' medical file.

Laboratory analyses

The levels of Malondialdehyde (MDA) and TAC in FF were considered as OS markers and have been assessed in Isfahan Cardiovascular Research Institute, Iran. Aliquots of the FF were thawed at room temperature and follicular fluid lipid peroxidation was assessed by MDA assay according to Oral et al (28). The TAC was assessed by enhanced chemiluminescence assay as described previously (29). All the samples were protected from direct sunlight.

Statistical analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences, version 13.00, SPSS Inc, Chicago, Illinois, USA (SPSS software). The normality of the vitamin intake level distributions was skewed. Therefore, normality of data was improved by using log transformation. Descriptive analyses were reported using mean and standard deviation. The data were analyzed using linear regression (adjusted for age, body mass index (BMI), smoking status,

calorie intake and etiology of infertility). Also, the daily vitamin intakes (vitamin A, C, and E) were divided into two groups according to the recommended dietary allowances and adequate intakes for adult females (USDA, Release 11, 1994) and the ART outcomes have been compared between the two groups using t-test. The P value of less than 0.05 was considered to be significant.

Results

In total, 219 subjects with mean age 31.54 ± 6.20 years participated in the study. The dietary vitamin intake levels in subjects and assisted reproduction outcomes were presented in table I.

Associations between the levels of vitamin intake and OS biomarkers in FF and ART parameters adjusted for age, physical activity, BMI, passive smoking status, etiology of infertility, and calorie intake are presented in table II.

The association between the vitamin intake levels and the MDA and TAC levels in FF were not significant.

Among the ART parameters, independent of the MDA and TAC levels in FF, the normal cleaved embryo rate was related to the vitamin A intake ($p < 0.05$); and the MII stage oocyte rate ($p = 0.02$), fertilization rate ($P < 0.05$) and percentage of good cleaved embryo ($p = 0.02$) were related to the vitamin C intake level positively.

In table III, the comparison of the OS biomarkers level in FF and ART parameters of 219 participants by the vitamins (vitamin A, C, and E) intake groups are shown. The mean of TAC level in FF among women who received vitamin C greater than 75 mg/d was higher than women with lower intakes.

The normal cleaved embryo rate was significantly higher in women with a vitamin A intake over 700 $\mu\text{g/d}$ than lower intakes. The MII stage oocyte rate, fertilization rate and percentage of non-segmented embryo were significantly higher in women with a vitamin C intake over than 75 mg/d than lower intakes.

Table I. Personal details, the level of the vitamin intake, and ART parameters in subjects

Variables	Mean±SD or %
Age (year)	31.54 ±5.45
BMI (kg/m ²)	26.60 ±4.33
Duration of infertility (year)	7.42±5.14
Etiology of infertility (%)	
Anovulation	40.64
Pelvic factor	46.58
Hypothalamic amenorrhea	5.48
Unexplained infertility	7.30
Vitamins intake level	
Vitamin A (µg/d)	1445.90 ±1684.50
Vitamin C (mg/d)	81.38±79.86
Vitamin E (mg/d)	22.84±18.45
Oxidative stress markers	
MDA (µ mol/lit)	0.98±0.29
TAC (molar Trolox equivalents)	1987.73±354.08
ART parameters	
Used gonadotropin (n)	40.54±1.72
Retrieved oocytes (n)	11.95±9.11
Fertilization rate (%)	62.60 ±27.10
Non-fragmented embryo rate (%)	44.81±36.21
Number of blastomers	6.13±2.44
Normal cleavage rate (%)	71.50±36.60
Good quality embryo rate (%)	47.14±34.01

BMI: Body mass index, MDA: Malondialdehyde, TAC: Total antioxidant capacity, ART: Assisted reproductive technology.

Table II. The correlation between dietary vitamin intake levels, oxidative stress markers and assisted reproduction parameters

	Number of oocytes			MII Rate			Fertilization rate			Normal cleaved embryo rate			Good embryo rate			Non- fragmented embryo rate		
	Beta	95% CI		Beta	95% CI		Beta	95% CI		Beta	95% CI		Beta	95% CI		Beta	95% CI	
Vitamin A (µg/d)	-0.11	-5.7	0.9	0.04	-8.5	9.5	-0.1	-0.2	0.2	0.2	0.1	12	0.07	-2.2	0.03	0.79	-9.2	16.1
Vitamin C (mg/d)	0.12	-0.6	5.3	0.2**	0.1	10	0.16*	0.01	0.3	0.2**	0.1	13.1	0.09	-5.4	18.9	0.02	-2.4	21.1
Vitamin E (mg/d)	-0.03	-5.4	3.6	-0.1	-17	5.4	0.02	-0.2	0.3	-0.1	-13	5.9	-0.07	-20	10.3	0.01	-10	10.2
MDA	-0.8	-0.7	2.2	-0.1	-1	10.7	-0.1	-0.4	0.1	-0.1	-33	0.14	-0.08	-31	9.2	-0.06	-18	13.8
TAC	-0.3	-0.01	0.003	0.06	-0.01	0.01	-0.1	-0.09	0.01	-0.02	-0.02	0.01	0.04	-0.01	0.02	0.08	-0.01	0.02

*p<0.05; **p=.02 (Multiple regression analysis)

CI: Confidence Interval, Beta: Standardized coefficient, MDA: Malondialdehyde, TAC: Total antioxidant capacity, MII: metaphase II.

Table III. Comparison of oxidative stress markers and ART parameters by dietary vitamin intake categorisations

	Vitamin A intake levels		p-value	Vitamin C intake levels		p-value	Vitamin E intake levels		p-value
	(µg/d)			(mg/day)			(mg/day)		
	≤700	>700		≤75	>75		≤15	>15	
Number	147	72		149	70		163	56	
MDA	0.99± 0.27	0.98± .31	0.55	0.97± 0.96	1.03± 0.82	0.17	0.99± 0.27	0.96± 0.33	0.21
TAC	1999.00± 345	1962± 375	0.21	1957.01± 380.02	2050.00± 283.11	0.04	1997.01± 343.11	1957± 387	0.15
No. of oocytes	12.11± 8.72	11.7± 9.60	0.19	11.81±9.03	12.90± 9.28	0.38	12.10± 9.41	11.6± 8.40	0.35
MIH Rate	82.00± 23.92	83.5± 15.51	0.20	79.01 ± 25.22	87.13± 14.81	0.008	83.24± 22.22	80.4± 23.70	0.20
Fertilization rate	65.62± 64.11	65.5± 63.41	0.28	60.31± 28.71	71.72± 22.41	0.04	64.9 ± 26.60	67.4± 29.30	0.09
Normal cleavage embryo rate	71.62± 36.21	81.9± 27.41	0.02	73.00± 35.31	80.5± 34.41	0.36	75.50± 34.23	73.5± 33.70	0.17
Good embryo rate	48.32± 36.90	52.11± 38.70	0.12	46.20± 36.51	55.80±34.40	0.18	49.13± 38.41	50.8± 35.10	0.36
Non-fragmented embryo rate	45.00± 35.32	50.2± 37.51	0.10	43.11± 35.42	56.21± 35.81	0.02	45.60± 35.81	49.92± 37.31	0.07

All data were presented as Mean±SD. ANOVA test.

ART: Assisted reproductive technology, MDA: Malondialdehyde, TAC: Total antioxidant capacity, MII: metaphase II.

Discussion

To determine the effect of dietary antioxidant vitamin intake on assisted reproduction treatment by influencing on the OS in follicular environment, two objectives were followed, including: the relations between the dietary vitamins A, C, and E intake levels and the measured the OS markers in FF and the relation between the dietary vitamin A, C and E intake and assisted reproductive parameters. Initially, the result of this study revealed that a relationship between intake of vitamins A, C and E intake and MDA and TAC in FF were not linear. However, in women with higher intakes of vitamin C level than the recommended levels (>75 mg/day),

the TAC level was significantly higher. Although, the effect of vitamins A, vitamin C and, vitamin E on maintaining the oxidant antioxidant balance in blood and tissues was demonstrated previously (16), these findings suggested that only dietary vitamin C intake may have a positive effect on antioxidant defense in follicular microenvironment.

Other findings revealed that the oocyte competence such as the oocyte maturation and fertilization were improved by more vitamin C intake, independent of the OS markers levels in FF. Also, higher dietary intake of this vitamin was associated with decreasing embryo fragmentation. The present observations were consistent with the previous studies showing protective effects of

vitamin C supplementation on MII mouse oocyte spindle structure and chromosomal alignment against an oxidative damage of hydrogen peroxide (30) and blocking the DNA damage in treating oocytes by ROS (31). Also, increased levels of glutathione peroxidase and vitamin C concentration, and reduced lipid peroxidation have been reported in FF of women who undergoing IVF program with multivitamin and mineral supplementation (18).

In some other reports, the beneficial effect of vitamin C on oocyte maturation was not supported. According to Dalvit et al, the use of the antioxidants vitamin E and C during in vitro maturation in bovine did not affect the percentage of nuclear maturation and fertilization (22). However, this study suggested that the assisted reproduction outcomes were not related to dietary intake level of vitamin E.

Therefore, the beneficial effect of vitamin C on ART outcome might not depend on the antioxidant effect of this vitamin. The effects of vitamin C on ART outcomes might be explained by anti-apoptotic effect of vitamin C on oocyte-granulosa cell complexes (32) and its role in regeneration of vitamin E has been demonstrated previously (33).

Also, this study revealed that although the association between the TAC level in FF and the vitamins A intake level was not significant, the percentage of normal cleaved embryo was positively related to vitamin A intake level. Also, these evidences suggested that the oocyte's developmental competence could be enhanced by vitamin A intake during intra follicular growth.

In consist with these findings, some researchers have suggested that antioxidants, including vitamins despite their antioxidant capacity might exert their effect on oocyte competence through other routes (18, 33). Combelles et al. suggested that in-vivo supplementation with vitamins that possess antioxidant activity did not necessarily improve reproduction (34).

The cytoplasmic retinol-binding proteins existed in human ovaries (35) and high concentration of vitamin A in human FF (36) suggested that vitamin A had an important role in reproductive function.

Conclusion

This paper described the impact of the dietary intake of vitamins on the oxidative protection in FF and on the oocytes-embryos parameters. The vitamins C and A might be useful to generate better eggs and embryos.

Acknowledgments

The authors appreciate Isfahan University of Medical Sciences and Tehran University of Medical Sciences for funding the survey.

Conflict of interest

Authors have no conflict of interests.

References

1. Combelles CM, Carabatsos MJ, Kumar TR, Matzuk MM, Albertini DF. Hormonal control of somatic cell oocyte interactions during ovarian follicle development. *Mol Reprod Dev* 2004; 69:347-355.
2. Sirard MA, Richard F, Blondin P, Robert C. Contribution of the oocyte to embryo quality. *Theriogenology* 2006; 65:126-136.
3. Hendriksen PJ, Vos PL, Steenweg WN, Bevers MM, Dieleman SJ. Bovine follicular development and its effect on the in vitro competence of oocytes. *Theriogenology* 2000; 53:11-20.
4. Bertout J, Mahutte NG, Preston SL, Behrman HR. Reactive oxygen species and ovarian function. In: Leung P; Adashi EY. *The Ovary*. San Diego: Elsevier Academic Press; 2004; 353-368.
5. Guerin P, El Mouatassim S, Menezo Y. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Hum Reprod Update* 2001; 7:175-189.
6. Matos L, Stevenson D, Gomes F, Silva-Carvalho JL, Almeida H. Superoxide dismutase expression in human cumulus oophorus cells. *Mol Hum Reprod* 2009;15:411-419.
7. Hanukoglu I. Antioxidant protective mechanisms against reactive oxygen species (ROS) generated by

- mitochondrial P450 systems in steroidogenic cells. *Drug Metab Rev* 2006; 38:171–196.
8. Zhang X, Li XH, Ma X, Wang ZH, Lu S, Guo YL. Redox-induced apoptosis of human oocytes in resting follicles in vitro. *J Soc Gynecol Investig* 2006; 13:451–458.
9. Valdez KE, Cuneo SP, Turzillo AM. Regulation of apoptosis in the atresia of dominant bovine follicles of the first follicular wave following ovulation. *Reproduction* 2005; 130: 71–81.
10. Benyamini Y, Gozlan M, Kokia E. Women's and men's perceptions of infertility and their associations with psychological adjustment: a dyadic approach. *Br J Health Psychol* 2009; 14:1-16.
11. Das S, Chattopadhyay R, Ghosh S, Goswami SK, Chakravarty BN, Chaudhury K. Reactive oxygen species level in follicular fluid embryo quality marker in IVF? *Hum Reprod* 2006; 21:2403–2407.
12. Oyawoye O, Abdel Gadir A, Garner A, Constantinovici N, Perrett C, Hardiman P. Antioxidants and reactive oxygen species in follicular fluid of women undergoing IVF: relationship to outcome. *Hum Reprod* 2003; 18: 2270-2274.
13. Song YL, Quan S, Tian JW, Li H, Chen SM, Xing FQ. Relationship between protein oxidation levels in the follicular fluid and the outcome parameters of in vitro fertilization-embryo transplantation. *Nan Fang Yi Ke Da Xue Xue Bao* 2009; 29:160-163.
14. Sefi M, Ben Amara I, Troudi A, Soudani N, Hakim A, Zeghal KM, et al. Effect of selenium on methimazole-induced liver damage and oxidative stress in adult rats and their offspring. *Toxicol Ind Health* 2012; 30:653-669.
15. Amin KA, Kamel HH, Abd Eltawab MA. Protective effect of Garcinia against renal oxidative stress and biomarkers induced by high fat and sucrose diet. *Lipids Health Dis* 2011; 10:6.
16. Sugino N. Reactive oxygen species in ovarian physiology. *Reprod Med Biol* 2005; 4: 31-44.
17. Tola EN, Mungan MT, Uğuz AC, Naziroğlu M. Intracellular Ca²⁺ and antioxidant values induced positive effect on fertilisation ratio and oocyte quality of granulosa cells in patients undergoing in vitro fertilisation. *Reprod Fertil Dev* 2013; 25:746-752.
18. Ozkaya MO, Naziroglu M. Multivitamin and mineral supplementation modulates oxidative stress and antioxidant vitamin levels in serum and follicular fluid of women undergoing in vitro fertilization. *Fertil Steril* 2010; 94: 2465-2466.
19. Arrigoni O, De Tullio MC. Ascorbic acid: much more than just an antioxidant. *Biochim Biophys Acta* 2002;1569:1-9.
20. Nadri B, Zeinoaldini S, Kohram H. Ascorbic acid effects on in vitro maturation of mouse oocyte with or without cumulus cell. *African Journal of Biotechnology* 2009; 8:5627-5631.
21. Thomas FH, Leask R, Srsen V, Riley SC, Spears N, Telfer EE. Effect of ascorbic acid on health and morphology of bovine preantral follicles during long-term culture. *Reproduction* 2001; 122: 487–495.
22. Dalvit G, Llanes SP, Descalzo A, Insani M, Beconi M, Cetica P. Effect of alpha-tocopherol and ascorbic acid on bovine oocyte in-vitro maturation. *Reprod Domest Anim* 2005; 40:93–97.
23. Duque P, Diez C, Royo L, Lorenzo PL, Carneiro G, Hidalgo CO, et al. Enhancement of developmental capacity of meiotically inhibited bovine oocytes by retinoic acid. *Hum Reprod* 2002; 17: 2706–2714.
24. Ikeda S, Kitagawa M, Imai H, Yamada M. The roles of vitamin A for cytoplasmic maturation of Bovine oocytes. *J Reprod Dev* 2005; 51: 23-35.
25. Laboratory manual of the WHO for the examination of human semen and sperm-cervical mucus interaction). *Ann Ist Super Sanita* 2001;37: 1-123.
26. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public Health Nutr* 2010; 13: 654-662.
27. Veeck LL. Oocyte assessment and biological performance. *Ann N Y Acad Sci* 1988; 541: 259-274.
28. Oral O, Kutlu T, Aksoy E, Fccolu C, Uslu H, Turul S. The effects of oxidative stress on outcomes of assisted reproductive techniques. *J Assist Reprod Genet* 2006; 23:81-85.
29. Attaran M, Pasqualotto E, Falcone T, Goldberg JM, Miller KF, Agarwal A, et al. The effect of follicular fluid reactive oxygen species on the outcome of in vitro fertilization. *Int J Fertil Women's Med* 2000;45:314-320.
30. Choi WJ, Banerjee J, Falcone T, Bena J, Agarwal A, Sharma RK. Oxidative stress and tumor necrosis factor-alpha-induced alterations in metaphase II mouse oocyte spindle structure. *Fertil Steril* 2007; 88:1220– 1231.
31. Tatemoto H, Ootaki K, Shigeta K, Muto N. Enhancement of developmental competence after in vitro fertilization of porcine oocytes by treatment with ascorbic acid 2-O-a-glucoside during in vitro maturation. *Biol Reprod* 2001; 65:1800–1806.
32. Eppig JJ, Hosoe M, OBrien MJ, Pendola FM, Requena A, Watanabe S. Conditions that affect acquisition of developmental competence by mouse oocytes in vitro: FSH, insulin, glucose and ascorbic acid. *Mol Cell Endocrinol* 2000; 163:109–116.
33. Botero D, Ebbeling CB, Blumberg JB, Ribaya-Mercado JD, Creager MA, Swain JF, et al. Acute effects of dietary glycemic index on antioxidant capacity in a nutrient-controlled feeding study. *Obesity* 2009;17:1664-1670.
34. Combelles CMH, Gupta S, Agarwal A. Could oxidative stress influence the in-vitro maturation of oocytes? *Reprod Biomed Online* 2009; 18: 864–880.
35. Ong DE, Page DL. Quantitation of cellular retinol-binding protein in human organs. *Am J Clin Nutr* 1986; 44: 425–430.

36. Schweigert FJ, Steinhagen B, Raila J, Siemann A, Peet D, Buscher U. Concentrations of carotenoids, retinol and alpha-tocopherol in plasma and follicular fluid of women undergoing IVF. *Hum Reprod* 2003; 18:1259–1264.