

# Potentials of the microalgae inoculant in restoration of biological soil crusts to combat desertification

A. Lababpour<sup>1</sup>

Received: 29 December 2015 / Revised: 1 July 2016 / Accepted: 12 July 2016 / Published online: 20 July 2016  
© Islamic Azad University (IAU) 2016

**Abstract** The world is presently faced to the many calamities, mainly the increased and rapidly developing environmental changes, soil degradation as an example of such environmental problems which is correlated with the destructive effects of the sandstorms. Biological soil crust (BSC), a main component of soil, has various environmental functions including reduction in the erosion by increasing soil stability and providing a sanctuary for the growth of the taxa and vascular plants communities. Destruction of BSC, which naturally can be recovered slowly in a long time processes, contributes to the desertification and other environmental catastrophes. Therefore, accelerating the BSC recovery both the quality and the quantity of the crust development, especially in the desert areas, is of the prime interest. Recent advances in the BSC restoration have provided an immense potential for emulating the natural restoration methods mainly through providing soils with inoculant. This paper reviews the present restoration-based procedures for the biological soil crust restoration practice. The main landmarks are presented and highlighted including strain(s) selection and development, mass biomass production, inoculum preparation, soil inoculation, soil augmentation, nurseries, and crust succession monitoring and control. The review also introduces several successful case studies in the USA and the Republic of China. Thereafter, the paper briefly documents the future directions of the research and technologies. Development of a restoration system through the

application of the microalgae inoculant is an encouraging aspect for accelerating the BSC recovery of the arid and semi-arid areas. However, further researches will help to establish and consolidate the potential of the microalgae cells and their application in desertification programs in large scales and in accordance with principles and requirements mandated by economic standards.

**Keywords** Biotechnological productivity · Erosion control · Microalgal mat · Soil community restoration · Soil health

## Introduction

Biological soil crust (BSC) has been the subject of interest of scientist from different science sectors, governments, communities, and international organizations such as United Nations (UNCCD 2015). The main reasons for such a high interest in BSC are the function, the vital role that it plays in the environment, along with the impacts that its disturbance would have on human life which threatens both the current, and the future lives on earth (UNCCD 2015). BSC is commonly defined as the outer cohesive thin horizontal ground cover, which is associated along with the living constituent that plays a vital ecological supporting roles such as protecting and stabilizing the soil against water and wind erosive forces, especially in the arid and semi-arid areas (Mor-Mussery et al. 2015).

Annually, a large scale and increasing amounts of environmental disturbances happen because of the direct or indirect disturbances in the global soil crust (Rosentreter et al. 2007). Several factors that cause BSC disturbances include climate change, soil degradation, desertification, vegetation depletion, drought, and low precipitation (i.e., rainfall) are among such factors. Loss of BSC and its

✉ A. Lababpour  
lababpour@niaeb.ac.ir

<sup>1</sup> Department of Industrial and Environmental Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran 14965-161, Iran



destruction results in loss of its ecological functions leading to the environmental consequences such as sand storms, reduction in soil stability, reduction in nitrogen and carbon fixation, water-related activities and effects on other soil microorganisms, and vascular plants will occur. The ultimate consequences of these destructive events may affect health, economic and other aspects of human life, or in the worse scenario, there may not be any chance of adjustment left forever (Field et al. 2010). Both the natural causes and human activities are the principal drivers of the processes of BSC degradation and desertification. In this regard, artificial restoration of the BSC may help to put an end and help the management of the desertification.

The various ecological functions of the BSC have been evidenced (Abdel-Raouf et al. 2012) through many functions that have been attributed to the BSC in practice including soil erosion reduction and control (Belnap et al. 2003). BSC performs its influence by aggregating unconsolidated soil particle both by the physical net and/or by releasing chemical substances such as polysaccharides (Abdel-Raouf et al. 2012). This function is partly executed by fixing the atmospheric carbon and nitrogen (Bu et al. 2013) and stabilization of the chemical elements such as chromium in the soil (Belnap et al. 2003), thereby influencing soil organic and inorganic nutrient materials (Belnap and Harper 1995). Further, BSC has a role in retaining soil moisture, so improving soil hydraulic conductivity (Belnap et al. 2003) and soil infiltration. BSC improves vegetation by helping germination of the seeds and inhibition of the weed, as well as protection against UV (Belnap et al. 2003; Lan et al. 2014a). BSC also has been used as an indicator of ecological integrity (Bowker et al. 2006).

The objective of this review article is to present a comprehensive review of the role of BSC and to provide a summary of the present methods of the BSC engineering, BSC restoration, and the challenges that might be encountered in this respect. The review mainly focuses on the crust microalgae as effective pioneer microorganisms in the restoration of the BSC. In addition, the main crust microalgae succession steps are explained and a number of successful implementation is introduced, accordingly. The article discusses the artificial BSC restoration following to an introduction, BSC definition, ecology, functions, roles, and the natural succession. This paper provides a critical review on microalgal-based BSC restoration. The main stages of strain(s) selection and development, mass production of microalgae, inoculant preparation, soil inoculation, soil augmentation, nurseries, and restoration process monitoring and controlling are highlighted. Challenges and future directions on BSC restoration with microalgae species are also outlined.

## Ecology of BSC

Often, the term BSC is referred to a consortium of autotrophic and heterotrophic living organisms in the outer layer of the soil surface. They may also be called cryptogamic, cryptobiotic (Clegg 2001), microbiotic, microphytic, and micropgytic soil crusts or soil- and rock-surface communities (SRSCs) (Pointing and Belnap 2012). Structurally, BSC consists of numerous kind of species including prokaryotic cyanobacteria (Brostoff et al. 2005), diatoms, eukaryotic microalgae (Cardon et al. 2008), lichen, mosses, fungi (Grishkan et al. 2015), liverworts, bacteria (Mengual et al. 2014a), and other taxa. BSC may be subdivided based on its biotic components such as cyanobacteria and microalgae crust (Karsten and Holzinger 2014a, b), lichen crust (Wu et al. 2011), mosses crusts, fungi crust, and bacterial crusts (Wang et al. 2007). Table 1 represents some of the genera and species of the BSC community. Geographically, BSC occurs in various areas and extreme habitats of the earth surface such as arid, hot, and cold desserts with a thickness ranging from few millimeters to several centimeters (Belnap et al. 2001). However, the community types and proportional amounts of the biological components of the crusts vary in different ecosystem and climates (Pietrasiak et al. 2013).

The natural sequences of BSC community succession generally include the development of cyanobacteria and microalgae, lichen and mosses, which is followed by other organisms and vegetation (Belnap et al. 2001). Figure 1 shows the stages of the BSC succession. Once a mat of these organisms has become established, the diversity of species increases through different succession stages and steps toward completion of the process (Bowker et al. 2006). Therefore, in developing engineered plans and methods for BSC restoration, all succession steps such as field practice should be considered to meet the desired crust formation.

## Conventional systems of BSC restoration

Several options are available to protect degraded crusts or accelerate the natural BSC restoration. Selection of the methods is mainly dependent on the location and the conditions of damaged crust. For example, in deserts with high speed winds, a frequently used conventional approach is wind shelter or check boards to provide protective conditions in the early crust development stages. In addition, land chemical mulching such as petrochemical is also used as pretreatments. Mulching inhibits soil erosion by providing a cover layer on the soil surface. However, the efficiency of the mulching is dependent on the mulch layer

**Table 1** Examples of the most common genera and species of the microorganisms in the natural BSC communities

Cyanobacteria and microalgae (Belnap 1995; Hu et al. 2002; Rahmonov and Piatek 2007; Karsten and Holzinger 2014b)	Lichen (Belnap 1995; Rosentreter et al. 2007; Wu et al. 2011)	Mosses (Rosentreter et al. 2007; Lan et al. 2012, 2014a; Xiao et al. 2015)
<i>Microcoleus vaginatus</i>	<i>Fulgensia fulgens</i>	<i>Bryum arcticum</i>
<i>Chroococcus minor</i>	<i>Psora decipiens</i>	<i>Didymodon vinealis</i>
<i>Chroococcus minutus</i>	<i>Squamarina</i> spp.	<i>Campylopus paradoxus</i>
<i>Chroococcus varius</i>	<i>Toninia sedifolia</i>	<i>Tortella tortuosa</i>
<i>Synechococcus aeruginosus</i>	<i>Catapyrenium</i> spp.	<i>Neckera crispa</i>
<i>Gloeocapsa atrata</i>	<i>Diploschistes</i> spp.	<i>Sphagnum</i>
<i>Merismopedia glauca</i>	<i>Endocarpon</i> spp.	<i>Tortula</i> spp.
<i>Microcoleus vaginatus</i> Gom	<i>Collema</i> spp.	<i>Bartramia pomiformis</i>
<i>Scytonema javanicum</i>		<i>Dicranum scoparium</i>
<i>Phormidium tenue</i>		<i>Bryum argenteum</i>
<i>Cylindrocapsa</i> sp.		
<i>Klebsormidium crenulatum</i>		
<i>Pinnularia borealis</i>		
<i>Stichococcus chlorelloides</i>		
<i>Stichococcus</i> cf. <i>fragilis</i>		


thickness and its resistance to erosion factors such as UV radiation, temperature, and characteristics of the soil particles (Chalker-Scott 2007). The combined systems of the mulching and windbreaks/shelterbelts have also been reported for reducing wind erosion in desert areas. Figure 2 shows an example of soil surface preparation before inoculation. However, these methods have temporary or insufficient effects and therefore should be renewed periodically. Vegetation is a useful environmental-friendly method and as a part of all features for combating desertification (Bainbridge and Darby 2014). However, in a study conducted by Lan et al., they compared the treatment of shifting sand dunes with *Salix (mongolica)* planting with and without inoculation of the cyanobacteria in an 8 years period and have found that only when *Salix* was planted to fix the sand dunes, the sand surface was relatively stabilized within a short time. When soil texture and nutrition had not been improved, only a number of vascular plants could survive at this time. However, after inoculation of sand dunes with the cyanobacteria, the BSCs restoration provided not only a more stable soil surface but also a favorable nutritional condition for the survival and succession of vascular vegetation communities. Therefore, compared to the shifting sand dunes and the dunes fixed with *Salix* planting, the higher vegetation coverage, biomass, and diversity could become attainable when dunes are covered with BSCs (Lan et al. 2014b).

The potential applications of the engineered systems for restoration of the existing BSC have a wide spectrum. Possibly the main advantage would be the acceleration in the succession procedure, a process that normally happens

very slowly in case of natural systems (Mor-Mussery et al. 2015). In addition, new restoration systems make the ability to design a real BSC for any required special restoration conditions possible (Büdel et al. 2014).

Generally, the process of natural BSC formation takes the following steps in a very long time (Campbell et al. 1989). In the primary step which normally happens in the first year, the sandy soil surface may become fixed by the bacteria adhering to the sand particles with exopolysaccharides (Paulo et al. 2012). Up to 4 years, the exposed crusts to the sandy surface would mainly be composed of the filamentous cyanobacterial communities that are dominated by the *Microcoleus* spp., which occurs as a cluster of filaments surrounded by a gelatinous sheath. The next step is the restoration of lichen and moss components. Establishment of the moss-dominated crusts is crucial for the establishment of the vegetation and restoration of the ecology (Rosentreter et al. 2007). The time course of each step may vary depending on the local situation and the biotic community such as annual rainfall, temperature, soil composition, and availability of autotrophic organisms. (Campbell et al. 1989). In addition, the natural BSC succession depends on the climate conditions, available species, as well as other ecological parameters. These parameters are out of control and therefore achievement of the desired BSC is almost impossible; however, the engineered BSC restoration may provide BSC design and implementation based on the site situations. This feasibility provides other alternatives for the BSC restoration in the arid and semi-arid areas as well as the subsequent vegetation with an accelerated process (Bowker 2007).






**Fig. 1** Succession stages of the BSC: **a** moving sand dune, **b** algae crust, **c** algae-lichen crust, **d** algae-moss crusts, **f**, **g** lichen-moss crusts, **h** moss crust (Lan et al. 2012), respectively







**Fig. 2** Generally, land preparation is required before soil inoculation, to provide inoculant succession. Picture shows land preparation by shelter or check boards in China (Hu et al. 2012)

### The general operation steps for the BSC restoration

The researches on the various BSC restoration concepts such as crust of microalgae, the crust of lichen, and the crust of mosses have been performed in the laboratory in addition to the field scale by various research groups (Xiao et al. 2015). These groups have often applied various techniques to their works in order to turn the unconsolidated sands into productive pasture or to accelerate the restoration of crust in the arid and semi-arid areas. The milestones in the restoration processes of the soil, as were identified and highlighted, might be divided and performed according to the following steps (Bu et al. 2013):

#### Criteria and restoration process design

Both biotic and abiotic criteria should be considered when an evaluation is taken under consideration for a BSC restoration plan. The main criteria in designing for a restoration plan includes definition of goals, determination of strategies, mode of operation (BSC restoration), thickness, crust development criteria, and distribution of BSC in the site (Liu et al. 2008). In addition, life cycle assessment and risk, possible combined options such as mulching, and implementation sequence stages are considered to be among these criteria. Other including factors are the required equipment and financial reserves, the ultimate site structure, capabilities and serviceability, and plans for monitoring and controlling including a list of present damaging factors, their influence and occurrence, and approaches for the modeling (Bowker et al. 2008). The main biotic parameters are present BSC community on the site, their pattern of distribution, required inoculant type and values, available options of inoculation, as well as feasible alternatives systems for massive production of the target biomass.

The main abiotic parameters are soil characteristics and topology, site climate condition including annual rainfall, temperature, humidity, and sunshine records as well as the possible available irrigation systems.

#### Preparation of the pure microalgae strain


The implementation process may start with strain preparation and development. Both pure microalgal culture and microalgae consortium of crust have been applied for BSC restoration in the laboratory and field studies (Lababpour et al. 2016). In some preliminary studies, the soil samples were harvested from a rich crust sites and inoculated into the new sites either directly or after dilution with water without further improving or cultivation in the open raceway or the closed photobioreactors (St. Clair and Johansen 1986). In the most published researches, mixed taxa were used for inoculation. For examples, moss components of the BSC have been used for BSC restoration by this method (Xiao et al. 2015). On the other hand, in other advanced technologies, the inoculant has been prepared by cultivation of the pure species in the open raceway systems and then was applied for inoculation (Liu et al. 2008). The strain preparation steps may include strain isolation and characterization by DNA extraction, amplification, and sequencing.

Figure 3 shows examples of the green algae isolated from desert habitats. In addition to taxonomic studies, diversity, physiology, and financial issues should also be subjected to the evaluations (Hu et al. 2012). Through, these evaluations and development of the methods were done along with other parameters including the effects of environmental parameters such as temperature, moisture, soil chemistry, and physical characteristics (Lin et al. 2013). In some studies, the applied microalgae strains were obtained from microalgae culture collections as a pure culture for BSC restoration studies (Rao et al. 2009).

#### Microalgae mass cultivation with high cell density

The selected species are mass cultivated in the open raceway or closed photobioreactors to obtain a large amount of the required biomass for inoculation. The open photobioreactors of up to 5 L bottles (Liu et al. 2008), 25L (Tan et al. 2012) and tank with the dimensions of  $2 \times 2 \times 1.5 \text{ m}^3$  (Wang et al. 2009) and large raceway (Rao et al. 2009) were reported for microalgae biomass production. Two main mass cultivation systems for the photosynthetic microorganisms are the open and closed photobioreactors (Pulz 2001). The open raceway cultivation system was reported for the inoculation of the microalgae crust in the greenhouse (Rao et al. 2009). However, the growth rate of microalgae is heavily





**Fig. 3** Examples of the common green algae isolated from desert habitats: **a** *Scenedesmus rotundus* (*Chlorophyceae*); **b** *Cylandrocystis* sp. (*Zygnematomyceae*); **c** *Myrmecia* sp. (*Trebouxiophyceae*); and

**d** *Chlorosarcinopsis* sp. (*Chlorophyceae*). Scale bar is equal to 10  $\mu$ m (Cardon et al. 2008)

**Table 2** Advantages and disadvantages of the liquid and solid inoculant preparation and field inoculation

	Advantages	disadvantages
Liquid inoculant	It is possible to include required micronutrients Preparation is easier compare to solid ones	Field inoculation timing is very important Storage is difficult Handling and transportation are difficult
Solid inoculant	Can be used in a loner time period compare to liquid ones Handling is easy	Field inoculation timing is not very important A carrier is required

dependent on the cultivation conditions. In addition, the mode of cultivation (autotrophic, heterotrophic, or mixotrophic) is options for cultivation which still remain to be developed for the crust restoration.

### Inoculant formulation and dosage

The produced biomass might directly be used as an inoculant in suspension or it might be used for further processing before it could be used (Sears and Prithiviraj 2012). Both of these systems have advantages and disadvantages. Table 2 lists the main advantages as well as disadvantages of the raw inoculant systems. The processes of dry inoculant preparation may include biomass harvesting, drying, and the addition of some additives to serve as a protector (Hu et al. 2012). Preparation of a large amount of the required inoculum with the desired quality and purity is attainable through the cultivation of the species; an approach that eliminates the need for extraction of the inoculums from the limited natural sources. Table 3 represents a summary of the features of the source of inoculants for biomass production and application to the soil.

### Soil inoculation

Both microalgal suspension and the dried microalgal biomass in the form of powder (normally a mixture of microalgae) have been applied as the soil inoculant (Hu et al. 2012). The microalgae inoculate contain viable cells and have the ability to germinate in the soil. The microalgae inoculant may be applied by tankers or agricultural aircraft (Hu et al. 2012; Sears and Prithiviraj 2012). In the case of direct application of the liquid inoculant, the inoculum concentration, timing, inoculum media, etc., should be considered before application in the field. The application of the liquid inoculant has been investigated in various concentrations and reported in the literature (Table 3). For example, a research group has used several concentrations of the microalgae inoculum and has found that 0.2 kg/ha of soil would be the best concentration of inoculation (Mengual et al. 2014b). Compare to dried inoculant, the direct application of the liquid inoculant does not need to a costly process of biomass harvesting, separation, and thickening, therefore, can be performed with less technological challenges and less expense.

On the other word, application of the dried biomass powder are easier in operation, simpler in transportations, and more stable for keeping for a long time. However, they are more expensive and their activity is always less effective compared to the liquid inoculants (Hu et al. 2012).

### Soil augmentation

In the field practical operations, the soil may be supplemented with the chemical nutrients (Liu et al. 2015). Using such fertilizers which support growth of inoculated viable cells depends on the soil characteristics as well as environmental climate conditions (Bertrand et al. 2014). In addition, the feasibility and availability of the fertilizers, as well as their cost, are important parameters that should be



**Table 3** Specifications of the applied liquid inoculant in the soil restoration programs so far

Inoculation method	Inoculated starter form	Value	References
Mosaic	Liquid as spray	0.3 mg Chl a/g dry soil	Liu et al. (2008)
Pure		1.6 g DCW/m <sup>2</sup>	Wang et al. (2009)
Mosaic	Spray	500 subsoil/500 mL H <sub>2</sub> O	St. Clair and Johansen (1986)
<i>Microcoleus vaginatus</i>	Sprinkling irrigation	1.0 mg in 2 × 15 m	Hu et al. (2002)

taken into the account when applications of the supplementary nutrients are desired (Khan et al. 2015). To improve the rate of BSC restoration in the field, recently biochar has been used in six concentration levels in a laboratory scale in order to investigate the effects of the supplementary biochar materials. Researchers have found that addition of 2 % biochar shows the highest influence on the microalgae concentration and the soil stabilization (Meng and Yuan 2014).

### Nurseries

To achieve a successful BSC restoration technology, it is necessary to know how various environmental factors influence BSC community at a site. A study on the effects of environmental factors: shade, water, and concentrations of the minerals such as nitrogen, phosphorous, potassium, and calcium during a 4-month period in Sonoran desert, USA, has resulted in the following order of effects:

$$\text{NH}_4\text{NO}_3 = \text{water frequency} > \text{shading} > \text{CaCO}_3 \\ = \text{KH}_2\text{PO}_4.$$

Soil water content was the primary positive factor (Bu et al. 2014). In addition, it is necessary to know how the inoculated communities interact with the biotic and abiotic factors and the best-matched strain for the specific condition. The interaction between soil cyanobacteria and the environmental parameters have been studied and reviewed over a long time period (Gómez et al. 2012; Raanan et al. 2015). From these results, it could be concluded that water availability and the related factors, such as frequency of rainfall, soil moisture content, together with the clay, and the amount of available phosphate in the soil, have the greatest influence on the cyanobacterial biomass concentration. Several other nursery challenges are soil cracking, crust, soil layers, thickness, cellular infiltration into the soil, adherence of the cells to the soil particles, etc.

### Monitoring and control of BSC dynamics

The monitoring of BSC development could be performed by a number of methods such as visual, light and electronic microscopic assays, molecular methods, or

estimated by computer simulations (Pelizer and Moraes 2014). Visual evaluation generally considers the color and morphology of the soil surface and compares it with the standard indexes (Belnap et al. 2008). In the other words, the analytical and molecular monitoring of the crust succession may include direct measurement of the pigments such as chlorophyll and molecular structure of the organisms. Measurement of the soil physical and chemical parameters is also common in the evaluation of BSC dynamics (Hu et al. 2003). For example, wind tunnel test is a method for evaluation of the soil stability which could be performed in the laboratory wind tunnels or in the field with a portable wind tunnel test apparatus (Zhao et al. 2010). The advanced remote crust chlorophyll monitoring system is an option that was reported for application in large areas of evaluation (Bu et al. 2013). A rational BSC monitoring will ensure a successful succession and reduction in the soil erodibility in the real field (Raggio et al. 2014).

### Successful case studies and challenges

Early BSC restoration practices in a laboratory scale were performed in the USA during the 1980s (St. Clair and Johansen 1986). Restoration was also practiced in the field in Colorado Plateau, Sonoran Desert, Mojave Desert, Chihuahuan Desert, Great Basin Desert, Florida shrubland, Massachusetts seashore, Oregon prairies, Wyoming steppe, Ohio, Michigan sand, and New Mexico (Bu et al. 2013). In the China, a widespread successful programs have been performed in the Kubuqi Desert, Hotq dessert, Loess plateau, Gurbantunggut Desert, Tengger Desert, Horqin Desert, Inner Mongolian steppe and Mu Us Desert during 2000s in the field area of up to 200 hectares supported by government and the United Nation (Bu et al. 2013). They have demonstrated a successful international cooperation to combat desertification. Results showed that cyanobacterial and algal soil coverage was increased up to 48.5 % and a total of 14 cyanobacterial and algal species were identified at the termination of inoculation experiment. The thickness, compressibility, and chlorophyll content of the biological crusts increased with inoculation time of 3–8 years; moss species appeared in



**Table 4** Examples of the commonly studied BSC sites and restoration around the world

Vicente Blanes Ecological Park in Molina de Segura	southeast Spain	Mengual et al. (2014b)
Vicente Blanes Ecological Park in Molina de Segura	Southeast Spain	Mengual et al. (2014a)
Vicente Blanes Ecological Park in Molina de Segura	Southeast Spain	Mengual et al. (2014b)
Vicente Blanes Ecological Park in Molina de Segura	Southeast Spain	Schoebitz et al. (2014)
Arches National Park	Moab, Utah, USA	Belnap and Harper (1995)
McMurdo Dry Valleys	South pole	McKnight et al. (2007)
Colorado Plateau region	South-eastern Utah, USA	Harper and Belnap (2001)
GSENM	Southern Utah, USA	Bowker et al. (2008)
Page	Arizona, USA	Bowker et al. (2008)
Naiman County, eastern part of Inner Mongolia	Republic of China	Zhao et al. (2010)
Arches National Park	Moab, Utah, USA	Belnap (1995)
Zhongwei County in the Ningxia Hui Autonomous Region	Republic of China	Grishkan et al. (2015)
southern part of the Pustynia Błędowska desert	Poland	Cabala and Rahmonov (2004)
Sahel area	Western Niger	Bertrand et al. (2014)
Western Mojave Desert	Los Angeles, CA	Brostoff et al. (2005)
western Negev Desert	Israel	Kidron et al. (2008)
Dugway Proving Grounds	Salt Lake City, Utah, USA	Belnap et al. (1993)
Coachella V alley	California, USA	Bainbridge and Darby (2014)
	South-eastern Utah, USA	Bowker et al. (2006)
Bocabec Bay	Southern New Brunswick, Canada	Tracy and South (1989)
Dalateqi county, Inner Mongolia	Republic of China	Wang et al. (2009)
Sonoran Desert near Phoenix	Arizona, USA	Bu et al. (2014)
Tengger Desert	Ningxia Hui Autonomous Region of China	Hu et al. (2002), (Liu et al. (2008)
	Southern Australia	Briggs and Morgan (2012)
Sonoran Desert	Southwestern United States	Bu et al. (2014)
Gurbantunggut Desert	Republic of China	Zheng et al. (2011)

the second year and cyanobacterial inoculation has resulted in an increased organic carbon in addition to the total nitrogen of the soil. The total salt, calcium carbonate, and electrical conductivity of the soil were also increased after inoculation. Diverse vascular plants communities such as *Artemisia ordosica* Krasch species were established after cyanobacterial inoculation on the windward and leeward slope surface of the dunes, respectively (Wang et al. 2009).


The American company, Soil Technologies Corp. has patented its technology and released its product into the market with a brand name of MICROP® (Soil Technologies Corp. 2015). The company uses the dried powder inoculant instead of the liquid suspension for soil inoculation. According to the company's report, the usefulness of the product was approved by the farmers that have used the product. In addition, TerraDerm Foundation has developed biological soil crust starter culture formulation technology with the trade names of TerraDerm and AgriDerm (Sears and Prithiviraj 2012). They have tested their products for soil stabilization and fertility in the USA and

Africa. Figure 4 shows the BSC restoration technology procedure presented by TerraDerm Foundation. They used agricultural aircraft for inoculation of the soil surface. They claimed successful BSC restoration in the used areas (Sears and Prithiviraj 2012). Table 4 presents several examples of the BSC practice sites around the world.

BSC succession is a long time dependent dynamic process in the field practice (Kumar and Adhikary 2015). In addition, the scale of world arid and semi-arid areas is 33.6–52.3 % of the terrestrial land surface and is growing fast (Reynolds et al. 2007). Therefore, the potential method should promote large landscape-scale applications and in an economical fashion. Hopefully, with the current rapid advancement in the microalgae-related technologies, the economic feasibility of the artificial BSC restoration systems is very close to the reality and attainable. Furthermore, any BSC restoration program requires management of the socioeconomic-related affairs of nearby communities which affect any restoration programs and should be considered before and during restoration program (Reynolds et al. 2007).







**Fig. 4** BSC succession stages as presented by the TerraDerm Foundation; **a** strain isolation from the natural sources; **b** strains purification and characterization stages; **c** mass cultivation of selected microorganisms in the artificial systems; **d** solid inoculant preparation

processes; **e** aerial soil inoculation; **f** irrigation; **g** deployment; **h** growth stage; **i** consortia emergence; and **j** vascular plant germination (Sears and Prithiviraj 2012)

## Conclusion and future directions

The main objective of the present review was to discuss the potential of the engineered BSC restoration technologies especially crust microalgae for recovery of the arid and semi-arid areas. These systems lead to an accelerated BSC recovery compared to the natural BSC long-term restoration systems. For large-scale restoration of BSC, mass production of microalgae in the open raceway or closed

photobioreactors systems could be a cost-effective option for providing large scale required inoculant. However, the efficiency of cultivation systems are still low and need further technological developments before they can be used in the large-scale field practices in the desertification programs.

Despite the technological challenges related to the cultivation, inoculation and nurseries, the potential benefits of the crust microalgae technologies are apparent.



Undoubtedly, implementation of the crust restoration in the arid and semi-arid areas presents major opportunities for the future research. Lastly, data on the effectiveness of the artificial BSC restoration in different soil types such as deserts, for instance, are needed to gain a realistic understanding of the impacts of microalgae strains on the BSC development and soil stabilization. In addition, the recent serious environmental and human health-related problems that are the consequence of the soil destruction are the leading causes and concerns for a drastic environmental change and its outcome such as sand-storm in dryland areas of the world, especially the Middle Eastern as well as African countries. This issue makes an urgent requirement for further researches and implementation of the BSC restoration projects, a subject which has been neglected so far.

**Acknowledgments** Support of the National Institute of Genetic Engineering and Biotechnology of IRAN is gratefully acknowledged (940801-I-535).

## References

- Abdel-Raouf N, Al-Homaidan AA, Ibraheem IBM (2012) Agricultural importance of algae. *Afr J Biotechnol* 11:11648–11658
- Bainbridge DA (2012) Restoration of arid and semi-arid lands. In: Aronson J, van Andel J (eds) *Restoration ecology: the new frontier*, 2nd edn. Wiley, Chichester
- Bainbridge DA, Darby MM (2014) Mitigation site soil characteristics and the effects of inoculation, nursery fertilizer practices and irrigation on survival and growth of mesquite (*Prosopis glandulosa*) in the Coachella Valley, California. *Restor Notes* 25:1–24
- Bashan Y, De-Bashan LE (2010) Microbial populations of arid lands and their potential for restoration of deserts. In: Patrice D (ed) *Soil biology and agriculture in the tropics*. Springer, Berlin, pp 109–137
- Belnap J (1995) Surface disturbances: their role in accelerating desertification. *Environ Monit Assess* 37:39–57. doi:10.1007/BF00546879
- Belnap J, Gillette DA (1997) Disturbance of biological soil crusts: impacts on potential wind erodibility of sandy desert soils in southeastern Utah. *L Degrad Dev* 8:355–362. doi:10.1002/(SICI)1099-145X(199712)8:4<355::AID-LDR266>3.0.CO;2-H
- Belnap J, Harper KT (1995) Influence of cryptobiotic soil crusts on elemental content of tissue of two desert seed plants. *Arid Soil Res Rehabil* 9:107–115. doi:10.1080/15324989509385879
- Belnap J, Harper K, Warren S (1993) Surface disturbance of cryptobiotic soil crusts: nitrogenase activity, chlorophyll content, and chlorophyll degradation. *Arid Land Res Manag* 8:1–8. doi:10.1080/15324989309381373
- Belnap J, Reynolds R, Reheis M, Phillips SL (2000) What makes the desert bloom? Contribution of dust and crusts to soil fertility on the Colorado Plateau. In: *ESA Annual Meeting*
- Belnap J, Kaltenecker JH, Rosentreter R et al (2001) Biological soil crusts: ecology and management: TR-1730-2. US Department of the Interior, Denver, CO
- Belnap J, Phillips SL, Duniway M, Reynolds RE (2003) Soil fertility in deserts: a review on the influence of biological soil crusts and the effect of soil surface disturbance on nutrient inputs and losses. In: Alsharhan AS, Wood WW, Goudie A et al (eds) *Desertification in the Third Millennium*. Swets & Zeitlinger, Lisse, pp 245–252
- Belnap J, Phillips SL, Witwicki DL, Miller ME (2008) Visually assessing the level of development and soil surface stability of cyanobacterially dominated biological soil crusts. *J Arid Environ* 72:1257–1264. doi:10.1016/j.jaridenv.2008.02.019
- Bertrand I, Ehrhardt F, Alavoine G et al (2014) Regulation of carbon and nitrogen exchange rates in biological soil crusts by intrinsic and land use factors in the Sahel area. *Soil Biol Biochem* 72:133–144. doi:10.1016/j.soilbio.2014.01.024
- Bowker MA (2007) Biological soil crust rehabilitation in theory and practice: an underexploited opportunity. *Restor Ecol* 15:13–23. doi:10.1111/j.1526-100X.2006.00185.x
- Bowker MA, Belnap J, Davidson DW, Goldstein H (2006) Correlates of biological soil crust abundance across a continuum of spatial scales: support for a hierarchical conceptual model. *J Appl Ecol* 43:152–163. doi:10.1111/j.1365-2664.2006.01122.x
- Bowker MA, Belnap J, Bala Chaudhary V, Johnson NC (2008) Revisiting classic water erosion models in drylands: the strong impact of biological soil crusts. *Soil Biol Biochem* 40:2309–2316. doi:10.1016/j.soilbio.2008.05.008
- Briggs AL, Morgan JW (2012) Post-cultivation recovery of biological soil crusts in semi-arid native grasslands, southern Australia. *J Arid Environ* 77:84–89. doi:10.1016/j.jaridenv.2011.10.002
- Brostoff WN, Rasoul Sharifi M, Rundel PW (2005) Photosynthesis of cryptobiotic soil crusts in a seasonally inundated system of pans and dunes in the western Mojave Desert, CA: field studies. *Flora Morphol Distrib Funct Ecol Plants* 200:592–600. doi:10.1016/j.flora.2005.06.008
- Bu C, Wu S, Xie Y, Zhang X (2013) The study of biological soil crusts: hotspots and prospects. *Clean Soil Air Water* 41:899–906
- Bu C, Wu S, Yang Y, Zheng M (2014) Identification of factors influencing the restoration of cyanobacteria-dominated biological soil crusts. *PLoS ONE* 9:e90049. doi:10.1371/journal.pone.0090049
- Büdel B (2003) Biological soil crusts of asia including the Don and Volga Region. In: Belnap J, Lange OL (eds) *Biological soil crusts: structure, function, and management ecological studies*. Springer, Berlin, pp 87–94
- Büdel B, Colesie C, Green TGA et al (2014) Improved appreciation of the functioning and importance of biological soil crusts in Europe: the Soil Crust International Project (SCIN). *Biodivers Conserv* 23:1639–1658. doi:10.1007/s10531-014-0645-2
- Cabala J, Rahmonov O (2004) Cyanophyta and algae as an important component of biological crust from the Pustynia Bledowska dessert (Poland). *Polish Bot J* 49:93–100
- Cabala J, Rahmonov O, Jablonska M, Teper E (2011) Soil algal colonization and its ecological role in an environment polluted by past Zn–Pb mining and smelting activity. *Water Air Soil Pollut* 215:339–348. doi:10.1007/s11270-010-0482-1
- Campbell SE, Seeler J, Golubic S (1989) Desert crust formation and soil stabilization. *Arid Soil Res Rehabil* 3:217–228
- Cardon ZG, Gray DW, Lewis LA (2008) The green algal underground: evolutionary secrets of desert cells. *Bioscience* 58:114. doi:10.1641/B580206
- Chalker-Scott L (2007) Impact of mulches on landscape plants and the environment—a review. *J Environ Hortic* 25:239–249
- Clegg JS (2001) Cryptobiosis—a peculiar state of biological organization. *Comp Biochem Physiol Part B Biochem Mol Biol* 128:613–624. doi:10.1016/S1096-4959(01)00300-1
- Durrell LW, Shields LM (1961) Characteristics of soil algae relating to crust formation. *Trans Am Microsc Soc* 80:73–79
- Field JP, Belnap J, Breshears DD et al (2010) The ecology of dust. *Front Ecol Environ* 8:423–430. doi:10.1890/090050



- Ge H, Xia L, Zhou X et al (2014) Effects of light intensity on components and topographical structures of extracellular polysaccharides from the cyanobacteria *Nostoc* sp. *J Microbiol* 52:179–183. doi:[10.1007/s12275-014-2720-5](https://doi.org/10.1007/s12275-014-2720-5)
- Gómez DA, Aranibar JN, Tabeni S et al (2012) Biological soil crust recovery after long-term grazing exclusion in the Monte Desert (Argentina). Changes in coverage, spatial distribution, and soil nitrogen. *Acta Oecol* 38:33–40. doi:[10.1016/j.actao.2011.09.001](https://doi.org/10.1016/j.actao.2011.09.001)
- Grishkan I, Jia R-L, Kidron GJ, Li X-R (2015) Cultivable microfungus communities inhabiting biological soil crusts in the Tengger Desert, China. *Pedosphere* 25:351–363
- Harper KT, Belnap J (2001) The influence of biological soil crusts on mineral uptake by associated vascular plants. *J Arid Environ* 47:347–357. doi:[10.1006/jare.2000.0713](https://doi.org/10.1006/jare.2000.0713)
- Hata T, Tsukamoto M, Inagaki Y et al (2011) Evaluation of multiple soil improvement techniques based on microbial functions. In: *Geo-Frontiers 2011*. American Society of Civil Engineers, Reston, pp 3945–3955
- Hu C, Liu Y, Song L, Zhang D (2002) Effect of desert soil algae on the stabilization of fine sands. *J Appl Phycol* 14:281–292. doi:[10.1023/A:1021128530086](https://doi.org/10.1023/A:1021128530086)
- Hu C, Zhang D, Huang Z, Liu Y (2003) The vertical microdistribution of cyanobacteria and green algae within desert crusts and the development of the algal crusts. *Plant Soil* 257:97–111. doi:[10.1023/A:1026253307432](https://doi.org/10.1023/A:1026253307432)
- Hu C, Gao K, Whitton BA (2012) Semi-arid regions and deserts. In: *Ecology of cyanobacteria II*. Springer, Dordrecht, pp 345–369
- Kabirov RR, Gaisina LA (2009) Parameters of the productivity of soil algae in terrestrial ecosystems. *Eurasian Soil Sci* 42:1374–1379. doi:[10.1134/S1064229309120072](https://doi.org/10.1134/S1064229309120072)
- Karsten U, Holzinger A (2014a) Green algae in alpine biological soil crust communities: acclimation strategies against ultraviolet radiation and dehydration. *Biodivers Conserv* 23:1845–1858. doi:[10.1007/s10531-014-0653-2](https://doi.org/10.1007/s10531-014-0653-2)
- Karsten U, Holzinger A (2014b) Green algae in alpine biological soil crust communities: acclimation strategies against ultraviolet radiation and dehydration. *Biodivers Conserv* 23:1845–1858
- Khan K, Pankaj U, Verma SK et al (2015) Bio-inoculants and vermicompost influence on yield, quality of *Andrographis paniculata*, and soil properties. *Ind Crops Prod* 70:404–409. doi:[10.1016/j.indcrop.2015.03.066](https://doi.org/10.1016/j.indcrop.2015.03.066)
- Kidron GJ, Vonshak A, Abeliovich A (2008) Recovery rates of microbiotic crusts within a dune ecosystem in the Negev Desert. *Geomorphology* 100:444–452. doi:[10.1016/j.geomorph.2008.01.012](https://doi.org/10.1016/j.geomorph.2008.01.012)
- Kumar D, Adhikary SP (2015) Diversity, molecular phylogeny, and metabolic activity of cyanobacteria in biological soil crusts from Santiniketan (India). *J Appl Phycol* 27:339–349. doi:[10.1007/s10811-014-0328-0](https://doi.org/10.1007/s10811-014-0328-0)
- Lababpour A, Kaviani M, Hosseini N (2016) Identification of prokaryotic microalgae from Asaluye area to improve soil and combat desertification. In: *1st International Conference on Dust*. Ahwaz, Iran
- Lan S, Wu L, Zhang D, Hu C (2012) Successional stages of biological soil crusts and their microstructure variability in Shapotou region (China). *Environ Earth Sci* 65:77–88. doi:[10.1007/s12665-011-1066-0](https://doi.org/10.1007/s12665-011-1066-0)
- Lan S, Wu L, Zhang D, Hu C (2013) Assessing level of development and successional stages in biological soil crusts with biological indicators. *Microb Ecol* 66:394–403. doi:[10.1007/s00248-013-0191-6](https://doi.org/10.1007/s00248-013-0191-6)
- Lan S, Wu L, Zhang D, Hu C (2014a) Desiccation provides photosynthetic protection for crust cyanobacteria *Microcoleus vaginatus* from high temperature. *Physiol Plant* 152:345–354. doi:[10.1111/ppl.12176](https://doi.org/10.1111/ppl.12176)
- Lan S, Zhang Q, Wu L et al (2014b) Artificially accelerating the reversal of desertification: cyanobacterial inoculation facilitates the succession of vegetation communities. *Environ Sci Technol* 48:307–315. doi:[10.1021/es403785j](https://doi.org/10.1021/es403785j)
- Li K, Bai Z, Zhang H (2014) Community succession of bacteria and eukaryotes in dune ecosystems of Gurbantünggüt Desert, Northwest China. *Extremophiles* 19:171–181. doi:[10.1007/s00792-014-0696-z](https://doi.org/10.1007/s00792-014-0696-z)
- Lin CS, Chou TL, Wu JT (2013) Biodiversity of soil algae in the farmlands of mid-Taiwan. *Bot Stud*. doi:[10.1186/1999-3110-54-41](https://doi.org/10.1186/1999-3110-54-41)
- Liu Y, Cockell CS, Wang G et al (2008) Control of Lunar and Martian dust—experimental insights from artificial and natural cyanobacterial and algal crusts in the desert of Inner Mongolia, China. *Astrobiology* 8:75–86. doi:[10.1089/ast.2007.0122](https://doi.org/10.1089/ast.2007.0122)
- Liu G, Yu H, Ma J et al (2015) Effects of straw incorporation along with microbial inoculant on methane and nitrous oxide emissions from rice fields. *Sci Total Environ* 518–519:209–216. doi:[10.1016/j.scitotenv.2015.02.028](https://doi.org/10.1016/j.scitotenv.2015.02.028)
- Makhalanyane TP, Valverde A, Gunnigle E et al (2015) Microbial ecology of hot desert edaphic systems. *FEMS Microbiol Rev* 39:203–221. doi:[10.1093/femsre/fuu011](https://doi.org/10.1093/femsre/fuu011)
- McKnight DM, Tate CM, Andrews ED et al (2007) Reactivation of a cryptobiotic stream ecosystem in the McMurdo Dry Valleys, Antarctica: a long-term geomorphological experiment. *Geomorphology* 89:186–204. doi:[10.1016/j.geomorph.2006.07.025](https://doi.org/10.1016/j.geomorph.2006.07.025)
- Meng X, Yuan W (2014) Can biochar couple with algae to deal with desertification? *J Sustain Bioenergy Syst* 04:194–198. doi:[10.4236/jsbs.2014.43018](https://doi.org/10.4236/jsbs.2014.43018)
- Mengual C, Roldán A, Caravaca F, Schoebitz M (2014a) Advantages of inoculation with immobilized rhizobacteria versus amendment with olive-mill waste in the afforestation of a semiarid area with *Pinus halepensis* Mill. *Ecol Eng* 73:1–8. doi:[10.1016/j.ecoleng.2014.09.007](https://doi.org/10.1016/j.ecoleng.2014.09.007)
- Mengual C, Schoebitz M, Azcón R, Roldán A (2014b) Microbial inoculants and organic amendment improves plant establishment and soil rehabilitation under semiarid conditions. *J Environ Manag* 134:1–7. doi:[10.1016/j.jenvman.2014.01.008](https://doi.org/10.1016/j.jenvman.2014.01.008)
- Meyer F, Bang S, Min S et al (2011) Microbiologically-induced soil stabilization: application of *Sporosarcina pasteurii* for fugitive dust control. *Geo-Frontiers 2011*:4002–4011
- Mor-Mussery A, Leu S, Budovsky A, Lensky I (2015) Plant-soil interactions and desertification: a case study in the northern Negev, Israel. *Arid Land Res Manag* 29:85–97. doi:[10.1080/15324982.2014.933455](https://doi.org/10.1080/15324982.2014.933455)
- Paulo EM, Boffo EF, Branco A et al (2012) Production, extraction and characterization of exopolysaccharides produced by the native *Leuconostoc pseudomesenteroides* R2 strain. *An Acad Bras Cienc* 84:495–508. doi:[10.1590/S0001-37652012000200018](https://doi.org/10.1590/S0001-37652012000200018)
- Pelizer LH, de Moraes IO (2014) A method to estimate the biomass of *Spirulina platensis* cultivated on a solid medium. *Braz J Microbiol* 45:933–936. doi:[10.1590/S1517-83822014000300024](https://doi.org/10.1590/S1517-83822014000300024)
- Pietrasiak N, Regus JU, Johansen JR et al (2013) Biological soil crust community types differ in key ecological functions. *Soil Biol Biochem* 65:168–171. doi:[10.1016/j.soilbio.2013.05.011](https://doi.org/10.1016/j.soilbio.2013.05.011)
- Pointing SB, Belnap J (2012) Microbial colonization and controls in dryland systems. *Nat Rev Microbiol* 10:551–562. doi:[10.1038/nrmicro2831](https://doi.org/10.1038/nrmicro2831)
- Pulz O (2001) Photobioreactors: production systems for phototrophic microorganisms. *Appl Microbiol Biotechnol* 57:287–293
- Raanan H, Oren N, Treves H et al (2015) Simulated soil crust conditions in a chamber system provide new insights on cyanobacterial acclimation to desiccation. *Environ Microbiol*. doi:[10.1111/1462-2920.12998](https://doi.org/10.1111/1462-2920.12998)



- Raggio J, Pintado A, Vivas M et al (2014) Continuous chlorophyll fluorescence, gas exchange and microclimate monitoring in a natural soil crust habitat in Tabernas badlands, Almería, Spain: progressing towards a model to understand productivity. *Biodivers Conserv* 23:1809–1826. doi:[10.1007/s10531-014-0692-8](https://doi.org/10.1007/s10531-014-0692-8)
- Rahmonov O, Piatek J (2007) Sand colonization and initiation of soil development by cyanobacteria and algae. *Ekológia (Bratislava)* 26:52
- Rao B-Q, Liu Y-D, Hu C-X et al (2009) The technology of man-made algal crust and its applications in control of desertification. *Acta Hydrobiol Sin* 33:756
- Reynolds JF, Smith DMS, Lambin EF et al (2007) Global desertification: building a science for dryland development. *Science* 316(80):847–851. doi:[10.1126/science.1131634](https://doi.org/10.1126/science.1131634)
- Rosentreter R, Bowker M, Belnap J (2007) A field guide to biological soil crusts of western U.S. drylands: common lichens and bryophytes. U.S. Government Printing, Denver
- Ruby-Pipeline-LLC (2012) Long-term monitoring plan for the Ruby Pipeline Project. [http://www.blm.gov/style/medialib/blm/nv/nepa/monitoring\\_reports/2012\\_year\\_1\\_monitoring.Par.54978.File.dat/Appendix\\_B\\_Long-Term\\_Monitoring\\_Plan.pdf](http://www.blm.gov/style/medialib/blm/nv/nepa/monitoring_reports/2012_year_1_monitoring.Par.54978.File.dat/Appendix_B_Long-Term_Monitoring_Plan.pdf)
- Schoebitz M, Mengual C, Roldán A (2014) Combined effects of clay immobilized *Azospirillum brasilense* and *Pantoea dispersa* and organic olive residue on plant performance and soil properties in the revegetation of a semiarid area. *Sci Total Environ* 466–467:67–73. doi:[10.1016/j.scitotenv.2013.07.012](https://doi.org/10.1016/j.scitotenv.2013.07.012)
- Sears JT, Prithviraj B (2012) Seeding of large areas with biological soil crust starter culture formulations: using an aircraft dispersible granulate to increase stability, fertility and CO<sub>2</sub> sequestration on a landscape scale. In: 2012 IEEE Green Technologies Conference. IEEE, pp 1–3
- Soil Technologies Corp. (2015) MICROP-SL. <https://www.soiltechcorp.com/search/results/?keywords=microp>. Accessed 27 July 2015
- St. Clair LL, Johansen JR (1986) Rapid stabilization of fire-disturbed sites using a soil crust slurry: inoculation studies. *Reclam Revegetation Res* 4:261–269
- Tan N, Jia SR, Han PP et al (2012) The open culture of *Nostoc flagelliforme* with a 25 L open pond. *Adv Mater Res* 554–556:1009–1012. doi:[10.4028/www.scientific.net/AMR.554-556.1009](https://doi.org/10.4028/www.scientific.net/AMR.554-556.1009)
- Tracy EJ, South GR (1989) Composition and seasonality of microalgal mats on a salt marsh in New Brunswick, Canada. *Br Phycol J* 24:285–291. doi:[10.1080/00071618900650301](https://doi.org/10.1080/00071618900650301)
- UNCCD (2015) Climate change and land degradation: bridging knowledge and stakeholders. In: UNCCD 3rd Scientific Conference. UNCCD, Cancun, Mexico
- Wang X, Chen F, Dong Z (2006) The relative role of climatic and human factors in desertification in semiarid China. *Glob Environ Chang* 16:48–57. doi:[10.1016/j.gloenvcha.2005.06.006](https://doi.org/10.1016/j.gloenvcha.2005.06.006)
- Wang W, Yang C, Tang D et al (2007) Effects of sand burial on biomass, chlorophyll fluorescence and extracellular polysaccharides of man-made cyanobacterial crusts under experimental conditions. *Sci China Ser C Life Sci* 50:530–534. doi:[10.1007/s11427-007-0051-z](https://doi.org/10.1007/s11427-007-0051-z)
- Wang W, Liu Y, Li D et al (2009) Feasibility of cyanobacterial inoculation for biological soil crusts formation in desert area. *Soil Biol Biochem* 41:926–929. doi:[10.1016/j.soilbio.2008.07.001](https://doi.org/10.1016/j.soilbio.2008.07.001)
- Wu L, Lan S, Zhang D, Hu C (2011) Small-scale vertical distribution of algae and structure of lichen soil crusts. *Microb Ecol* 62:715–724. doi:[10.1007/s00248-011-9828-5](https://doi.org/10.1007/s00248-011-9828-5)
- Xiao B, Zhao Y, Wang Q, Li C (2015) Development of artificial moss-dominated biological soil crusts and their effects on runoff and soil water content in a semi-arid environment. *J Arid Environ* 117:75–83. doi:[10.1016/j.jaridenv.2015.02.017](https://doi.org/10.1016/j.jaridenv.2015.02.017)
- Xie Z, Liu Y, Hu C et al (2007) Relationships between the biomass of algal crusts in fields and their compressive strength. *Soil Biol Biochem* 39:567–572. doi:[10.1016/j.soilbio.2006.09.004](https://doi.org/10.1016/j.soilbio.2006.09.004)
- Zhao H-L, Guo Y-R, Zhou R-L, Drake S (2010) Biological soil crust and surface soil properties in different vegetation types of Horqin Sand Land, China. *CATENA* 82:70–76. doi:[10.1016/j.catena.2010.05.002](https://doi.org/10.1016/j.catena.2010.05.002)
- Zheng Y, Xu M, Zhao J et al (2011) Effects of inoculated *Microcoleus vaginatus* on the structure and function of biological soil crusts of desert. *Biol Fertil Soils* 47:473–480. doi:[10.1007/s00374-010-0521-5](https://doi.org/10.1007/s00374-010-0521-5)

