

ANALYSIS OF THE EXTRACTS OF *ISATIS TINCTORIA* BY NEW ANALYTICAL APPROACHES OF HPLC, MS AND NMRJue Zhou, Ph.D.,^{1*} Fan Qu, Ph.D.,²

¹ College of Food Science and Biotechnology, Zhejiang Gongshang University, Hangzhou, Zhejiang, China, ² Women's Hospital, School of Medicine, Zhejiang University, Hangzhou, Zhejiang, China, ³ Pharmaceutical Sciences Division, King's College London, Franklin-Wilkins Building, 150 Stamford Street, London SE1 9NH, UK

*Email: juezhou2006@yahoo.com.cn

Abstract

The methods of extraction, separation and analysis of alkaloids and indole glucosinolates (GLs) of *Isatis tinctoria* were reviewed. Different analytical approaches such as High-pressure Liquid Chromatography (HPLC), Liquid Chromatography with Electrospray Ionization Mass Spectrometry (LC/ESI/MS), Electrospray Ionization Time-Of-Flight Mass Spectrometry (ESI-TOF-MS), and Nuclear Magnetic Resonance (NMR) were used to validate and identity of these constituents. These methods provide rapid separation, identification and quantitative measurements of alkaloids and GLs of *Isatis tinctoria*. By connection with different detectors to HPLC such as PDA, ELSD, ESI- and APCI-MS in positive and negative ion modes, complicated compounds could be detected with at least two independent detection modes. The molecular formula can be derived in a second step of ESI-TOF-MS data. But for some constituents, UV and MS cannot provide sufficient structure identification. After peak purification, NMR by semi-preparative HPLC can be used as a complementary method.

Keywords: *Isatis tinctoria*; HPLC; MS; NMR; indole glucosinolates (GLs); alkaloids.

Introduction

Isatis tinctoria is widely used as anti-inflammatory and dye medicinal plant in Europe and China. It has been used and cultivated in Europe and China since antiquity. Over the last decade, more than 65 compounds of *Isatis tinctoria* including glycosides, alkaloids (Wu et al, 1997; Liu et al, 2001a,b; Liu et al, 2003), organic acids (Wang et al, 2009), and sucrose (Liu et al, 2002a,b) were identified. *Isatis tinctoria* contains indolic compounds. Among these compounds, glucobrassicin and its derivatives have antitumoral effect, especially against mammary cancer (Hoessel et al, 1999; Xu et al, 1991; Lin et al, 2002; Thornalley 2002; Galletti et al, 2006).

Different analytical approaches such as High-Pressure Liquid Chromatography (HPLC) (Qin et al, 2001; Tu et al, 2009), Liquid Chromatography with Electrospray Ionization Mass Spectrometry (LC/ESI/MS), Electrospray Ionization Time-Of-Flight Mass Spectrometry (ESI-TOF-MS), and Nuclear Magnetic Resonance (NMR) were used for analysis of the compounds in *Isatis tinctoria*. These methods provide rapid separation, identification and quantitative measurements of glucosinolates and alkaloids from *Isatis tinctoria*. The methods of extraction, preparation and analysis of glucosinolates and

alkaloids of *Isatis tinctoria* were reviewed in this paper.

Alkaloids

Indigo is one of the oldest natural blue dye (Uzal et al, 2010). Indole is a product of tryptophan catabolism by gut bacteria and is absorbed into the human body in substantial amounts (Gillam et al, 2000). Indole is absorbed and metabolized within the liver to indican (indol-3-ylsulfate) (Fordtran et al, 1964; Levy 1995). Indigo and indirubin have been found in human urine. Indigo and indirubin are structural isomers, which have physiological effects on liver microsomes in mice (Sugihara et al, 2004). Tryptanthrin, is an indoloquinazoline alkaloid that strongly inhibits cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5-LOX) in cellular assays (Henning et al, 2002a; Henning et al, 2002b; Meragelman et al, 2002; Chan et al, 2009).




Figure 1: Relationship of tryptophan, indigo, indirubin and indican

Extraction and separation

Freshly harvested materials of *Isatis indigotica* were cut into small pieces of 2–3 cm length and immediately shock frozen with liquid nitrogen. Prior to extraction, the leaves were dried at 40 °C. Temperature, relative humidity and weight loss was monitored during the drying process. Constant weight was achieved after 3–4 days. The dried leaf samples were stored at room temperature for a few days in brown glass bottles in the dark. Immediately before extraction, dried leaf material was grounded. 1.0 g frozen and powdered samples was extracted by Pressurised Liquid Extraction (PLE) with an ASE 200 instrument (Liau et al, 2007). Bagci and Özçelik (2009) investigated fatty acid in the seeds of *Isatis tinctoria* using GC and HPLC.

Sample preparation of Mass Spectrometry

Standard solutions (0.1 mg/ml) of indigo, isoindigo, indirubin, isatin, indican, indoxyl acetate and 2-indolinone were prepared by dissolving 5 mg of each powder in 50 ml of DMSO. The solutions were filtered over a 0.45 µm syringe filter. The solution of the mixture of active agents was prepared daily by dilution of appropriate volumes of the standard solutions with acetonitrile.

New analytical approaches

The methods used to identify indigos and its isomers include spectrophotometer in the UV, visible and IR regions (Cordy and Yeh, 1984; Miliani et al, 1998; Vautier et al, 2001), micro-Raman spectroscopy (Vandenabeele and Moens, 2003), electron spin resonance (Russell and Kaupp, 1969), thin layer chromatography (Kokubun et al, 1998) and liquid chromatography with UV/visible detection (Wouters and Rosario-Chirinos, 1992; Cooksey and Tyrian, 2001; Maugard et al, 2001; Orska-Gawry's et al, 2003). The alkaloids 5-hydroxyoxindole (5), 3-(20-carboxyphenyl)-quinazolin-4-on (8) and bisindigotin (21) have been described for *Isatis tinctoria* or *Isatis indigotica* before (Wu et al, 1997; Li et al, 2000; Wei et al, 2005). The available reference compounds were identified by NMR spectra. To identify indigos and its isomers, compounds were identified on the basis of their UV data in combination with MS, high resolution-MS, and literature data on their chromatographic retention (Canjura and Schwartz, 1991; van Breemen et al, 1991; Terasaki et al, 2002; de Rosso and Mercadante, 2007a, b). HPLC profiles obtained with PDA, ELSD, MS and MS/MS detectors are given as supplementary data (Van Breemen et al, 1991). APCI-MS and MS/MS data were compared with the published literature (de Rosso and Mercadante, 2007a, b). The parameters of selected indigoid compounds and their precursors are presented in Table 1.

Table 1: The parameters of selected indigoid compounds and their precursors

Class	Compounds	λ_{max} (nm) (DMSO)	$[M + H]^+$	Reference
Indigoid coloring agents	Indigo, indigotin(blue)	617	263	Puchalska et al, 2004
	Indirubin (red)	540	263	
	Isoindirubin (red)	552	263	
	Isoindigo (brown)	365, 490	263	
Indigoid precursors	Indican	218, 280	296	
	Isatan	218, 280	310	
	Indoxyl acetate	280	176	
	2-Indolinone	-	134	
	Isatin	416	148	

HPLC method was widely used to analyze alkaloids in medicinal plants. The parameters of tryptanthrin, indigo, and indirubin analysis by HPLC are presented in Table 2.

Based on thin layer chromatography (TLC) and HPLC with an ultraviolet detector (HPLC-UV) (Ding et al, 2001; Henning et al, 2002a), LC/ESI/MS and fast atom bombardment mass spectrometry (FAB-MS) has been used to analyze *Isatis tinctoria* (Wouters et al, 1991; Ozeki et al, 1993; Sun et al, 2000). Recently, scientists have used LC/MS to investigate indirubin and indigo in textiles and historical art objects (Szostek et al, 2003; Puchalska et al, 2004). Liau et al used LC/APCI/MS to quantify the amounts of tryptanthrin, indigo, and indirubin in *Isatis tinctoria* (Zhang 1983; Liau et al, 2007).

Mass spectrometry (MS) has been applied to the identification of indigoid compounds using direct inlet into different ion sources: electron ionization (EI) (McGovern et al, 1991; Clark and Cooksey, 1997) laser desorption/ionization (LDI) and APCI (Szostek et al, 2003; Bagci and Özçelik, 2009). The previous study reported LC-APCI-MS method is developed for detecting and analyzing tryptanthrin, indigo, and indirubin in the leaves and roots of *Isatis indigotica*

(Oberthür et al, 2003; Liao et al, 2007).

Table 2: The parameters of tryptanthrin, indigo, and indirubin by HPLC

Compounds	Alkaloids : tryptanthrin, indigo, and indirubin
HPLC	¹ 1100 liquid chromatography system equipped with two pumps, a MWD UV detector and Rheodyne injector. ² L-4250 UV-VIS detector and an L-6200A pump
Column	¹ Thermo Hypurity-Advence C18 (5µm, 250mm×3mm i.d.) (USA) column and a Phenomenex Luna Security Guard Cartridge C18 (5µm, 4mm×2.0mmi.d.) ² ZORBAX ODS
Mobile phases	A : water 0.005% trifluoroacetic acid (TFA); B: ACN containing 0.005% TFA
Conditions	¹ Gradient for the separation of fraction 2 was: 5% B for 2 min, 5–65% B in 98 min, 65–70% B in 10 min; gradient for the separation of fraction 3 was: 34% B for 13 min, 34–85% B in 87 min. 210 nm ² B: 0-5-30 min; 20-50-100% at a flow rate of 2 ml/min;552nm
Reference	¹ Liao et al, 2007 ² Aobchey et al, 2007

Reversed-phase LC/ESI/MS and UV/visible spectrophotometric methods were elaborated for the identification of indigoid (indigo, indirubin, isoindigo, isoindirubin) color components of natural dye stuffs and their natural or synthetic precursors (indican, isatin, indoxyl, 2-indolinone) (Puchalska et al, 2004).

Gillam et al (2000) reported the definitive identification of indigo and indirubin as products of human cytochrome P450 (P450)-catalyzed metabolism of indole by visible, ¹H NMR and MS. Mohn et al (2009) discovered a new indolic alkaloid from *Isatis tinctoria* by combined several detectors such as photodiode array detector (PDA), ESI/MS, and APCI-MS (positive and negative ion modes). Structural information was obtained by unspecific evaporative light scattering detector (ELSD)/MS/MS (Xiao et al, 2007) experiments and by high-resolution mass spectra recorded by ESI-TOF-MS.

Different analytic methods have been used to identify and determine the compounds in *Isatis tinctoria*. Previous reports reveal that LC/MS does not require sample pretreatment and allows for the detection of tryptanthrin, indigo, and indirubin at the same time. ESI/MS with high sensitivity, offers detection limits in the range 0.03–5.00 µg/ml for these compounds examined. Some selected parameters of these methods used to analyze alkaloid in *Isatis tinctoria* are presented in Table 3.

Previous reports show the concentrations (µg/g) of tryptanthrin, indigo, and indirubin in different samples from *Isatis indigotica* as follows: tryptanthrin 0.110, 0.614, 0.533, 0.409, 0.164, 0.153; Indigo 0.016, 1.859, 1.685, 1.562, 0.693, 0.271; Indirubin 0.224, 2.384, 1.148 (You et al, 1998; Liang et al, 2000; Liao et al, 2007). Puchalska (2004) reported the results of indigoids by HPLC/ESI/MS as presented in Table 4.

Indole glucosinolates (GLs)

Extraction and separation

Indole glucosinolates (GLs) are highly reactive compounds, with a varying aglycon side chain. According to the type of this chain they are classified into aliphatic, aromatic and indole GLs. Among the latter, glucobrassicin (GBS), together with other indol-type GLs, was isolated from *Isatis tinctoria* (Elliott and Stowe, 1970; Elliott and Stowe 1971;

Table 3: The parameters of tryptanthrin, indigo, and indirubin analyzed by MS

Products	Alkaloids : tryptanthrin, indigo, and indirubin
Column	Zorbax SB-C18 (150 d 4.6 mm, 3.5 μ m)
Instrument	HPLC: ¹ LC: Surveyor liquid chromatography system. MS: APCI, quadrupole ion trap instrument. ² 1100 LC MS: High-Resolution ESI/TOF /MS (BioTOF III; Bruker Daltonics, Inc. Billerica, MA, USA)
Eluent	A, ACN; B, 0.15% formic acid
Gradient program	A: 3 min 20% 8 min 50% 10 min 55% 22 min 55% 35 min 100% 45 min 100% Flow rate:0.6 ml/min; Injection volume: 5 μ l; wavelength: 280, 610 nm
MS parameters	Sheath gas for tryptanthrin, indigo or indirubin. aux gas 1.5 L/min for tryptanthrin, Nebulizing gas flow and temperature: 10 ml/min and 300 °C Vaporization temperature: 575 °C; capillary temperature, 200 °C
Voltage	Discharge: 3.5 μ A for tryptanthrin indigo, and 5.5 μ A for IS. capillary voltage, 10V. Ionization voltage: 4000 V (positive ion mode); Orifice voltage: 90 V
Instrument	LC-MSD 1100 (Agilent Technologies)
m/z	Observed mass range (TIC): 50–500 Observed selected ions (SIM): 134, 148, 176, 205, 263, 296, 338, 340, 418, 420, 422
References	Liau et al, 2007 Chanayath et al, 2002; Puchalska et al, 2004

Table 4: The results of indigoids by HPLC/ESI/MS

Parameter	Indigotin	Isoindigo	Indirubin	Isatin	Reference
Selected ion, <i>m/z</i>	263	263	263	148	Puchalska et al, 2004
Calibration curve $y = ax + b$					
a	598367	341891	319965	80950	
b	16205	430	47560	27036	
Linearity range (µg/ml)	0.03–4.20	0.03–4.00	0.03–5.00	0.45–18.20	
Detection limit(µg/ml)	0.01	0.01	0.01	0.15	
Quantification limit(µg/ml)	0.03	0.03	0.03	0.45	

Frechard et al, 2001). Some reports demonstrated that glucosinolates is very sensitive to temperature. 60–70 °C have been used in validated PLE methods for many classes of natural products such as isatin A and B (Benthin et al., 1999; Oberthür et al, 2003; Oberthür et al, 2004a, b; Basalo et al, 2006). For the extraction and isolation of GLs of *Isatis tinctoria*, existing methods (Kushad et al, 1999; Kiddle et al, 2001; Verkerk et al, 2001; Bennett et al, 2007) has a problem with the extent of thermal degradation at temperature above 50 °C with. Extractions were typically carried out at temperatures between 70 °C and 100 °C. Thermal degradation of GLs during cooking has been reported (Slominski and Campbell, 1989; Oerlemans et al, 2005; Bones and Rossiter, 2006).

Mohn et al (2007a) isolated GLs from *Isatis tinctoria* seeds by Soxhlet-extraction for 8 h with 400mL petrol ether (boiling range 40–60 °C). After evaporation to dryness, the residue was extracted three times with water (room temperature, 3×150 mL), centrifuged (5 min, 1600×*g*, room temperature) and filtered. The aqueous solutions were concentrated to 45mL in vacuo, and 5mL acetonitrile were added. After centrifugation, the supernatant was introduced into a column packed with DEAE-Sephadex A-25 (50 g). The column was eluted with water/acetonitrile 80:20 until the eluate was colorless. GLs were eluted with a mixture of 0.1M K₂SO₄/acetonitrile 80:20 at 3.2 mL/min and monitored with UV detection at 229 nm. Fractions were analyzed by HPLC–MS. Finally, isolated GLs were freeze dried. Purity and structures of isolated compounds were confirmed by NMR and LC–MS experiments.

New analytical approaches

In the recent years, new analytical approaches such as LC-based on-line spectroscopy, MS, NMR opened new avenues for increasingly comprehensive analysis of plant extracts and secondary metabolites. Researches by LC-based on-line spectroscopy has been widely taken on metabolite profiling studies (Burns et al, 2003; Yamazaki et al, 2003; Le Gall et al, 2005; Long et al, 2006; Dan et al, 2008; Ding et al, 2008; Iijima et al, 2008; Qiao et al, 2008; Schliemann et al, 2008). Previous reports on the indigo precursors and GLs showed harvest regimen affected the chemical composition of *Isatis tinctoria* leaves (Oberthür et al, 2004b; Mohn and Hamburger, 2008). Mohn et al (2008) carried out a quantitative assay for the direct analysis of intact GLs in *Isatis tinctoria* leaves. In their research, pressurized liquid extraction (PLE) was used. Detection was carried out by ESI/MS in the negative ion mode. Mohn and Hamburger (2008) used atmospheric pressure chemical ionization (APCI) and ESI/MS, and ESI-TOF-MS detectors to detect the extracts of *Isatis tinctoria*.

The dichloromethane extracts of *Isatis tinctoria* are lipophilic with active pharmacological effect (Oberthür, 2003). Mohn et al (2009) used dichloromethane with PLE method to extract lipophilic compounds. The parameters of the

dichloromethane extracts by HPLC are presented in table 5.

Table 5: The parameters of the dichloromethane extracts by HPLC

Extracts	The dichloromethane extracts
HPLC	Agilent series 1100 system equipped with degasser, binary high pressure mixing pump, column thermostat and photodiode array (PDA) detector
Column	SunFire C18 column, 5 μ m, 150 $\times$ 10.0 mm I.D., equipped with a guard column (10.0 $\times$ 10.0 mm I.D)
Mobile phases	A: water with formic acid 0.1%, B: acetonitrile, flow rate: 0.5 ml/min,
Conditions	5% B isocratic for 2 min, 5%–93% B in 30 min, 93% B isocratic for 17 min, 93–100% B in 6 min. column temperature:25.0 $^{\circ}$ C,
Reference	Mohn et al, 2009

In previous reports, trifluoroacetic acid and formic acid are often used as mobile phase for chromatography of intact GLs (Mellon et al., 2002; Bringmann et al., 2005; Song et al., 2005). The parameters in Table 6 present GLs extracts analysed by HPLC-PDA-MS and HPLC-TOF-MS. The parameters of LC/TOF/ MS methods for GLs are presented in Table 7.

Table 6: The parameters of the dichloromethane extracts analyzed by LC/ESI /MS

Extracts	The dichloromethane extracts
Instrument	¹ The second splitter outlet: evaporative light scattering detector (ELSD series 2000, Alltech, Deerfield IL, USA) APCI (Bruker Daltonics; Bremen, Germany) Esquire 3000 plus ion trap ESI /MS
Conditions	¹ Ion charge conditions (positive mode: ICC 30000; negative mode: ICC 20000), scan speed:13000 m/z/s, gauss filter: 0.2 m/z, trap drive: 58.8 (negative mode: 61.4), isolation width: 4.0 m/z, fragmentation amplitude: 1.00 V. ESI: A:B= 1:3 Nitrogen pressure =2.0 bar Reference solution of sodium formate 0.1% in isopropanol/water (1:1) containing 5 mM sodium hydroxide.
Voltage	Capillary voltage:-4500 V (negative mode: +4500 V), endplate offset: -500 V, capillary exit voltage: 128.5 V (negative mode: -128.5 V), skimmer voltage: 40 V (negative mode:-40 V) hexapole at 250.0 Vpp (negative mode:230 Vpp) skimmer 1 at 40 V (negative mode: -50 V) skimmer 2 at 22.5 V (negative mode: -22.5 V).
Gas	Drying gas: nitrogen, flow rate 5 μ l/min

	nebulising gas (pressure: 30 psi), collision gas: helium
Temperature	55 °C, nebuliser temperature:375 °C, Drying temperature: 250 °C. Dry gas temperature 240 °C.
m/z	100–800 in positive and negative modes for the detection of carotenoids and porphyrins in the hexane/acetone extract: 100–1200
References	Mohn et al, 2009

Table 7: The parameters of GLs analyzed by LC/TOF/ MS

Products	GLs: sinigrin, epiprogoitrin, progoitrin, gluconapin, glucotropaeolin, glucobrassicin, neoglucobrassicin, sulfolucobrassicin, 4-hydroxyglucobrassicin
Instrument	LC/TOF/ MS, Negative ion LC–MS spectra on the ion trap
Voltage	Capillary voltage was at 4500V, endplate offset at –500V, capillary end voltage at –115.0V, skimmer voltage:–40.0V and trap drive at 53.4.
Gas	drying gas: nitrogen flow rate of 10 L/min, nebulizing gas: pressure of 30 psi
Temperature	nebulizer temperature : 300 °C
m/z	100–800 negative mode:100–600
Instrument calibration	a reference solution of sodium formate 0.1% in isopropanol/water (1:1) containing 5mM sodium hydroxide.
References	Mohn et al, 2009

Some minor constituents purified by semi-preparative HPLC were analyzed by off-line microprobe NMR spectroscopy. The analysis of GLs by NMR are presented in Table 8.

Table 8: The parameters of GLs analyzed by NMR

Products	NMR	References
GLs	NMR spectra were obtained with a 500 MHz Avance III system equipped with a 1 mm TXI probe. Data processing was with Topspin 2.1 Bruker DRX500 NMR spectrometer equipped with a 5mm SEI probe I.S.: 1,3,5-trimethoxybenzene	Mohn et al, 2009 Mohn et al, 2007a,b,c

Isatis tinctoria has been cultivated throughout Europe, especially in Western and southern Europe, since ancient times. In China, *Isatis tinctoria* has been used thousands of years as an important and popular herbal medicine in Traditional Chinese Medicine (TCM) for the treatment of inflammatory diseases. This paper reviewed the extraction methods of hydrophilic compounds (alkaloids) and lipophilic compounds (indole glucosinolates) of *Isatis tinctoria*. Multiple analytical approaches including HPLC, LC/ESI/MS, ESI-TOF-MS, and NMR have been reported as validation and quality control

tools of this herbal medicine. HPLC separates complicated compounds. By connection with different detectors to HPLC such as PDA, ELSD, ESI- and APCI-MS in positive and negative ion modes, complicated compounds could be detected with at least two independent detection modes. The molecular formula can be derived in a second step of ESI-TOF-MS data. But for some constituents, UV and MS cannot provide sufficient structure identification. After peak purification, NMR by semi-preparative HPLC can be used as a complementary method. Selected parameters of these analytical methods are summarized in this article. This review may be useful to guide the analysis of the extracts of *Isatis tinctoria* in the future.

Acknowledgements

Dr. Fan QU was funded by National Natural Science Foundation of China (No.30800390), China Postdoctoral Science Foundation Key Project (No. 200902642), China Postdoctoral Science Foundation (No.20070421188), Outstanding Young Medical Scientist Foundation of Zhejiang Province (No. 2008QN022) and Zhejiang Traditional Chinese Medicine Foundation (No. 2008YB010).

References

1. Aobchey, P., Sinchaikul, S., Phutrakul, S. and Chen, S.T. (2007). Simple Purification of Indirubin from *Indigofera tinctoria* Linn. and Inhibitory Effect on MCF-7 Human Breast Cancer Cells. *Chiang Mai J. Sci.* **34**: 329-337
2. Bağcı, E. and Özçelik, H., (2009). Fatty acid and tocopherol patterns of some *Isatis L.* (brassicaceae) species from turkey. *Pak. J. Bot.*, 41: 639-646.
3. Basalo, C., Mohn, T., and Hamburger, M., (2006). Are extraction methods in quantitative assays of pharmacopoeia monographs exhaustive? A comparison with pressurized liquid extraction. *Planta Med.* **72**: 1157.
4. Bennett, R.N., Carvalho, R. Mellon, F.A., Eagles, J., and Rosa, E.A.S, (2007). Identification and Quantification of Glucosinolates in Sprouts Derived from Seeds of Wild *Eruca sativa L.* (Salad Rocket) and *Diplotaxis tenuifolia L.* (Wild Rocket) from Diverse Geographical Locations. *J. Agric. Food Chem.* **55**: 67.
5. Benthin, B., Danz, H., and Hamburger, M. (1999). Efficient method for preparation of highly purified lipopolysaccharides by hydrophobic interaction chromatography. *J. Chromatogr. A* **732**:39-41.
6. Bones, A.M., and Rossiter, J.T., (2006). The enzymic and chemically induced decomposition of glucosinolates. *Phytochemistry* **67**: 1053-1064.
7. Bringmann, G., Kajahn, I., Neusuess, C., Pelzing, M., Laug, S., Unger, M., and Holzgrabe, U. (2005). Analysis of the glucosinolate pattern of *Arabidopsis thaliana* seeds by capillary electrophoresis coupled to electrospray ionization time-of-flight mass spectrometry. *Electrophoresis.* **26**: 1513-1522.
8. Burns, J., Fraser, P.D., and Bramley, P.F., (2003). Identification and quantification of carotenoids, tocopherols and chlorophylls in commonly consumed fruits and vegetables. *Phytochemistry* **62**: 939-947.
9. Canjura, F.L., and Schwartz, S.J., (1991). Separation of chlorophyll compounds and their polar derivatives by high-performance liquid chromatography. *J. Agric. Food Chem.* **39**: 1102-1105.
10. Chan, H.L., Yip, H.Y., Mak N.K., and Leung, K.N. (2009). Modulatory Effects and Action Mechanisms of Trypantanthrin on Murine Myeloid Leukemia Cells. *Cellular & Molecular Immunology*, 6 :335-342
11. Chanayath, N., Lhieochaiphant, S. and Phutrakul, S., (2002). Pigment extraction techniques from the leaves of *Indigofera tinctoria* Linn. and *Baphicacanthus cusia* Brem. and chemical structure analysis of their major components. *CMU J.*, **1**: 149-160.

12. Clark, R.J.H, and Cooksey, C.J. (1997). Bromindirubins: the synthesis and properties of minor components of Tyrian Purple and the composition from *Nucella lapillus*. J. Soc. Dyers Colour. 113: 316.
13. Cooksey, C., and Tyrian, J. (2001). Purple: 6, 60-dibromoindigo and related compounds. Molecules **6**: 736.
14. Cordy, A., and Yeh, K., (1984). Blue dye identification on cellulosic fibers: indigo, logwood, and Prussian blue. J. Am. Instit. Conserv. **24**: 33.
15. Dan, M., Su, M., Gao, X., Zhao, T., Zhao, A., Xie, G., Qiu, Y., Zhou, M., Liu, Z., and Jia, W., (2008). Metabolite profiling of *Panax notoginseng* using UPLC-ESI-MS. Phytochemistry **69**: 2237–2244.
16. de Rosso, V.V., and Mercadante, A.Z., (2007a). HPLC-PDA-MS/MS of anthocyanins and carotenoids from dovyalis and tamarillo fruits. J. Agric. Food Chem. **55**: 9135–9141.
17. de Rosso, V.V., and Mercadante, A.Z., (2007b). Identification and quantification of carotenoids, by HPLC-PDA-MS/MS, from Amazonian fruits. J. Agric. Food Chem. **55**: 5062–5072.
18. Ding, S. P., Liu, Y. H., Li, J., Qin, G. W. and Wu, X. Y. (2001). Studies on chemical constituents of *Isatis indigotica* Fort. (Cruciferea) II. China J. Herald of Med. **20**: 475-476.
19. Ding, S., Dudley, E., Plummer, S., Tang, J., Newton, R.P., and Brenton, A.G, (2008). Fingerprint profile of Ginkgo biloba nutritional supplements by LC/ESI-MS/MS. Phytochemistry **69**: 1555–1564.
20. Elliott, M.C. and Stowe, B.B. (1970). A novel sulphonated natural indole. Phytochemistry **9**:1629–1632
21. Elliott, M.C. and Stowe, B.B. (1971). Distribution and variation of indole glucosinolates in woad (*Isatis tinctoria* L.). Plant Physiol **48**:498–503
22. Fordtran, J. S., Scroggie, W. B., and Polter, D. E. (1964). Colonic absorption of tryptophan metabolites in man. J. Lab. Clin. Med. **64**, 125-132.
23. Frechard, A., Fabre, N., Pean, C., Montaut, S., Fauvel, M-T., and Rollin, P., (2001). Novel indole-type glucosinolates from woad (*Isatis tinctoria* L.). Tetrahedron Lett **42**:9015–9017
24. Galletti S., Barillari J., Iori R. and Venturi G. (2006). Glucobrassicin enhancement in woad (*Isatis tinctoria*) leaves by chemical and physical treatments. Journal of the Science of Food and Agriculture. **86**:1833–1838
25. Gillam E.M. J., Notley L.M., Cai H., De Voss J.J., and Guengerich F. P. (2000). Oxidation of indole by cytochrome P450 enzymes. Biochemistry **39**, 13817-13824
26. Henning D., Dietmar, B., Matthias, H. (2002a). Inhibitory activity of tryptanthrin on prostaglandin and leukotriene synthesis. Planta Med. **68** 875–880.
27. Henning D., Stefka, S., Petra, W., Axel, B., and Matthias, H. (2002b). Quantitative determination of the dual COX-2/5-LOX inhibitor tryptanthrin in *Isatis tinctoria* by ESI-LC-MS. Planta medica, **68**:152–157.
28. Hoessel, R., Leclerc, S., Endicott, J.A., Nobel, M.E., Lawrie, A., Tunnah, P., Leost, M., and Damiens E, (1999). The active constituent of a Chinese antileukaemia medicine, inhibits cyclindependent kinases. Nature Cell Biol. **1**: 60.
29. Iijima, Y., Suda, K., Suzuki, T., Aoki, K., and Shibata, D., (2008). Metabolite profiling of chalcones and flavanones in tomato fruit. J. Jpn. Soc. Hort. Sci. **77**, 94–102.
30. Kiddle G., Bennett R.N., Botting N.P., Davidson N.E., Robertson A.A.B. and Wallsgrove R.M., (2001). High-performance liquid chromatographic separation of natural and synthetic desulphoglucosinolates and their chemical validation by UV, NMR and chemical ionisation-MS methods. Phytochem. Anal. **12**: 226-242.
31. Kokubun, T, Edmonds, J, and John, P. (1998). Indoxyl derivatives in woad in relation to medieval indigo production. Phytochemistry, **49**: 79.
32. Kushad, M.M., Brown A.F., Kurilich A.C., Juvik J.A., Klein B.P., Wallig M.A., and Jeffery E.H. (1999). Variation of Glucosinolates in Vegetable Crops of Brassica oleracea. J. Agric. Food Chem. **47**: 1541.
33. Le Gall, G., Metzдорff, S.B., Pedersen, J., Bennett, R.N., Colquhoun, I.J., (2005). Metabolite profiling of *Arabidopsis thaliana* (L.) plants transformed with an antisense chalcone synthase gene. Metabolomics **1**: 181–198.

34. Levy, H.L. (1995). Hartnup Disorder. In *Metabolic and Molecular Bases of Inherited Disease*, CR Scriver, AL Beaudet, WS Sly, D Valle, eds (New York: McGraw-Hill), pp 3629–3642.
35. Li, B., Chen, W.S., Zheng, S.Q., Yang, G.J., and Qiao, C.Z., (2000). Two new alkaloids isolated from tetraploid Banlangen. *Yao Xue Xue Bao* **35**: 508–510.
36. Liang, Y. H. , Hou, H. X. , Li, D. R. , Qin, Q. , Qiu, L. and Wu, H. H. (2000). Studies on *in vitro* anticancer activity of tryptanthrin B. *China J. Chin. Tradit. Herbal Drugs* **31**: 531-533.
37. Liao, B.C, Jong, T.T., Lee, M.R., Chen, S.S., (2007). LC-APCI-MS method for detection and analysis of tryptanthrin, indigo, and indirubin in Daqingye and Banlangen. *Journal of pharmaceutical and Biomedical Analysis*. **43** : 346-351
38. Lin, A. H. , Fang, S. X. , Fang, J. G , Du, G. and Liu, Y. H. (2002) Studies on anti-endotoxin activity of F022 from radix isatidis. *China J. Chin. Materia Medica* **27**: 439-442
39. Liu, Y. H., Duing, S. P., Lin, A. H., Fang, J. G. and Shi, S. P. (2001a). The protective effects of different polar fractions from radix isatidis on endotoxin challenged mice. *China J. Acta. Univ. Med. Tongji* **30**: 272-273.
40. Liu, Y. H., Qin, G. W., Ding, S.P. and Wu, X.Y. (2002b) Studies on chemical constituents in Root of *Isatis indigotica* III. *China J. Chin. Tradit. Herbal Drugs* **33**: 97-99.
41. Liu, Y. H., Qin, G.W. , Ding, S.P. and Wu, X.Y. (2001b). Studies on chemical constituents in root of *Isatis indigotica* I. *China J. Chin. Tradit. Herbal Drugs* **32**: 1057-1060.
42. Liu, Y.H., Qin, G.W., Fang, J. G. and Wu, X. Y. (2002a). Screening of chemical constituents with antiendotoxin activity from radix isatidis. *Herald of Med.* **21**: 74-75.
43. Liu, Y.H., Wu, X.Y., Fang, J.G., and Tang , J., (2003). Studies on Chemical Constituents from Radix Isatidis(IV). *Herald of Med* **9**: 591-594.
44. Long, M., Millar, D.J., Kimura, Y., Donovan, G., Rees, J., Fraser, P.D., Bramley, P.M., and Bolwell, G.P., (2006). Metabolite profiling of carotenoid and phenolic pathways in mutant and transgenic lines of tomato: identification of a high antioxidant fruit line. *Phytochemistry* **67**: 1750–1757.
45. Maugard, T, Enaud, E., Choisy, P., and Legoy, M.D. (2001). Identification of an indigo precursor from leaves of *Isatis tinctoria* (woad). *Phytochemistry* **58**: 897.
46. McGovern, P.E., Lazar, J., and Michel, R.H. (1991). Caveats on the analysis of indigoid dyes by mass spectrometry. *J. Soc. Dyers Colour.* **107**: 280.
47. Mellon, F.A., Bennett, R.N., Holst, B., and Williamson G. (2002). Intact Glucosinolate Analysis in Plant Extracts by Programmed Cone Voltage Electrospray LC/MS: Performance and Comparison with LC/MS/MS Methods. *Analytical Biochemistry* **306**: 83–91.
48. Meragelman, K.M., West, L.M., Northcote, P.T., Pannell, L. K., McKee, T.C., and Boyd, M.R. (2002). Unusual Sulfamate Indoles and a Novel Indolo [3,2-a] carbazole from *Ancorina* sp. *J. Org. Chem.* **67**: 6671-6677.
49. Miliani, C., Romani, A., and Favaro, G. (1998). A spectrophotometric and fluorimetric study of some anthrax quinoid and indigoid colorants used in artistic paintings. *Spectrochim. Acta, Part A.* **54**: 581.
50. Mohn, T., Plitzko I., Hamburger M. (2009). A comprehensive metabolite profiling of *Isatis tinctoria* leaf extracts. *Phytochemistry*, **70**: 924–934.
51. Mohn, T., Cutting, B., Ernst, B., Hamburger, M., (2007a). Extraction and analysis of intact glucosinolates – a validated pressurized liquid extraction/liquid chromatography–mass spectrometry protocol for *Isatis tinctoria*, and qualitative analysis of other cruciferous plants. *J. Chromatogr. A* **1166**: 142-151.
52. Mohn, T., Hamburger, M., (2008). Glucosinolate pattern in *Isatis tinctoria* and *I. Indigotica* seeds. *Planta Med.* **74**: 885–888.
53. Mohn, T., Potterat, O., and Hamburger, M. (2007b). A comprehensive metabolite profiling of *Isatis tinctoria* leaf

- extracts. *Phytochemistry*. **70**: 924-934.
54. Mohn, T., Potterat, O., and Hamburger, M., (2007c). Pigments in leaf extracts of *Isatis tinctoria* by HPLC/UV/MS. *Planta Med.* **73**: 151–156.
 55. Mohn, T., Suter, K., and Hamburger, M. (2008). Seasonal changes and effect of harvest on glucosinolates in *Isatis* leaves. *Planta Med.* **74**: 582–587.
 56. Oberthür C., Graf H., and Hamburger M. (2004a). Determination of indican, isatin, indirubin and indigotin in *Isatis indigotica* by liquid chromatography/electrospray ionization tandem mass spectrometry. *Phytochemistry*, **65**: 3261.
 57. Oberthür, C., Graf, H., Hamburger, M., (2004b). The content of indigo precursors in *Isatis tinctoria* leaves - a comparative study of selected accessions and postharvest treatments. *Phytochemistry*, **65**: 3261–3268.
 58. Oberthür, C., Heinemann C., Elsner, P., Benfeldt E., and Hamburger M., (2003). A comparative study on the skin penetration of pure tryptanthrin and tryptanthrin in *Isatis tinctoria* extract by dermal microdialysis coupled with isotope dilution ESI-LC-MS. *Planta Med.* **69**: 385–389.
 59. Orska-Gawry's, J., Surowiec, I., Kehl, J., Rejniak, H., Urbaniak-Walczak, K., and Trojanowicz, M. (2003). Identification of natural dyes in archeological Coptic textiles by liquid chromatography with diode array detection. *J. Chromatogr. A*. **989**: 239.
 60. Ozeki, N., Oka, H., Ikai, Y., Ohno, T., Hayakawa, J., Sato, T., Ito, M., and Suzuki, R. (1993). Identification of coal-tar dyes by reversed-phase TLC/FABMS with a sample condensation technique. *J. Food Hyg. Soc. Japan*, **34**: 512–516.
 61. Puchalska, M., Poleć-Pawlak, K., Zadrozna, I., Hryszko, H. and Jarosz, M., (2004). Identification of indigoid dyes in natural organic pigments used in historical art objects by high-performance liquid chromatography coupled to electrospray ionization mass spectrometry. *J. Mass Spectrom.* **39**: 1441–1449.
 62. Qiao, X., Han, J., Xu, M., Zhan, Z., Sun, S., Li, H., Wang, B., Ye, M., and Guo, D. (2008). Characterisation of chemical constituents in Guan Xin II decoction by liquid chromatography coupled with electrospray ionization-mass spectrometry. *Planta Med.* **74**: 1720–1729.
 63. Qin, Q., He, H.P., Soren, B.C., Hasse, B.R., Arsalan, K. and Ming, C. (2001). Isolation of low polar fractions from banlangen and its immunocompetence. *Chin. J. Clin. Pharm.* **10**: 29-31.
 64. Russell, G.A., and Kaupp, G. (1969). Oxidation of carbanions. IV. Oxidation of indoxyl to indigo in basic solution. *J. Am. Chem. Soc.* **91**: 3851.
 65. Schliemann, W., Ammer, C., and Strack, D. (2008). Metabolite profiling of mycorrhizal roots of *Medicago truncatula*. *Phytochemistry*, **69**: 112–146.
 66. Slominski, B. A., and Campbell, L. D. (1989). Gas chromatographic determination of indole glucosinolates — A re-examination. *J. Sci. Food Agric.* **40**: 131–143.
 67. Song, L., Morrison, J.J., Botting, N.P., and Thornalley, P.J. (2005). Analysis of glucosinolates, isothiocyanates, and amine degradation products in vegetable extracts and blood plasma by LC–MS/MS. *Anal. Biochem.* **347**: 234-243.
 68. Sugihara, K., Kitamura, S., Yamada, T., Okayama, T., Ohta, S., Yamashita, K., and Matsuda, T. (2004). Aryl hydrocarbon receptor-mediated induction of microsomal drug-metabolizing enzyme activity by indirubin and indigo. *Biochem Biophys Res Commun.* **318**: 571-578.
 69. Sun, G.L., Hu, Z.L., Meng, H., Li, Y., Wang, X. L. and Liu, J. H. (2000). Detection of anti- cytomegalovirus effect of radix isatidis by MTT method. *Shan Dong Univ. of TCM.* **24**: 137-138.
 70. Szostek, B., Orska-Gawry's, J., Surowiec, I., Trojanowicz, M. (2003). Investigation of natural dyes occurring in historical Coptic textiles by high-performance liquid chromatography with UV–Vis and mass spectrometric detection. *J. Chromatogr. A* **1012**: 179.
 71. Terasaki, M., Itabashi, Y., Suzuki, T., and Nishimura, K., (2002). An improved method for determining the composition of FFA in red tide flagellates by RP-HPLC with fluorescence detection. *J. Am. Oil Chem. Soc.* **79**:

- 1181–1186.
72. Thornalley, P.J., (2002). Isothiocyanates: Mechanism of cancer chemopreventive action. *Anti-Cancer Drugs*, 13: 331
 73. Tu, Y.Q., Sun, J., Ge, X., Li, Z.Y. (2009). Chromosome elimination, addition and introgression in intertribal partial hybrids between *Brassica rapa* and *Isatis indigotica*. *Annals of Botany*. 103:1039-1048.
 74. Uzal, N., Yilmaz, L., Yetis, U. (2010). Separation Science and Technology, Nanofiltration and Reverse Osmosis for Reuse of Indigo Dye Rinsing Waters. *Separation Science and Technology*. 45:331 - 338
 75. Van Breemen, R.B., Canjura, F.L., Schwartz, S.J., (1991). Identification of chlorophyll derivatives by mass spectrometry. *J. Agric. Food Chem.* **39**: 1452–1456.
 76. Vandenabeele, P., Moens, L. (2003). Micro-Raman spectroscopy of natural and synthetic indigo samples. *Analyst*. 128: 187.
 77. Vautier, M., Guillard, C., and Herrmann, J.M. (2001). Photocatalytic degradation of dyes in water: case study of indigo and of indigo carmine. *J. Catal.* **201**: 46.
 78. Verkerk, R., Dekker, M., and Jongen, W.M.F.(2001). Post-harvest increase of indolyl glucosinolates in response to chopping and storage of Brassica vegetables. *J. Sci. Food Agric.* **81**: 953.
 79. Wang, Y., Qin, Q., Chen, J., Kuang, X., Xia, J., Xie, B., Wang, F., Liang, H. and Qi, Z.Q. (2009). Synergistic effects of *Isatis tinctoria* L. and tacrolimus in the prevention of acute heart rejection in mice. *Transplant Immunology*. **22**: 5-11
 80. Wei, X.Y., Leung, C.Y., Wong, C.K.C., Shen, X.L., Wong, R.N.S., Cai, Z.W., and Mak, N.K., (2005). Bisindigotin, a TCDD antagonist from the Chinese medicinal herb *Isatis indigotica*. *J. Nat. Prod.* **68**: 427–429.
 81. Wouters, J. and Verhecken, A., (1991). High-performance liquid chromatography of blue and purple indigoid natural dyes. *J. Soc. Dyers Colour.*, 107: 266-269
 82. Wouters, J., Rosario-Chirinos, N. (1992). Dye analysis of pre-Columbian Peruvian textiles with high-performance liquid chromatography and diode array detection. *J. Am. Inst. Conserv.* 31: 237.
 83. Wu, X. Y. , Liu, Y. H. , Sheng, W. Y. , Sun, J. and Qin, G. W. (1997). Chemical constituents of *Isatis indigotica*. *J. Planta Med.* **63**: 55-57.
 84. Xiao, S.S., Kaishun B., and Sun, Y.Q., (2007). Identification of Chemical Constituents in the Root of *Isatis Indigotica* Fort. by LC/DAD/ESI/MS/MS. *J. Liquid Chromatography & Related Technologies*, **30**: 73 - 85
 85. Xu, Y. M. , Lu, P. C. , Wang, Y. Z. and Huang, Z. X. (1991). Research in amylose of *Isatis indigotica* Fort. for enhancing immunity. *Chin. J. of Integrated Tradit. and Western Med.* 11: 357-359.
 86. Yamazaki, Y., Urano, A., Sudo, H., Kitajima, M., Takayama, H., Yamazaki, M., Aimi, N., and Saito, K., (2003). Metabolite profiling of alkaloids and strictosidine synthase activity in camptothecin producing plants. *Phytochemistry* **62**: 461–470.
 87. You, S. , Yao, X. S. , Chen, Y. J. , Cui, C. B. , Qu, G. X. , Wu, J. and Xu, X. J. (1998). Research in effective anticoagulant components of *Isatis indigotica* Fort. *China J. Bull. Chin. Materia Medica* **13**: 31-32.
 88. Zhang, S. X. (1983). Studies on chemical constituents root of *Isatis indigotica*. *China J. Chin. Tradit. Herbal Drugs* 14: 7-9.